



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ : C12N 15/54, 15/82, 15/11, 9/10, 5/10, C08B 30/00, A01H 5/00, A23L 1/0522	A1	(11) International Publication Number: WO 97/45545 (43) International Publication Date: 4 December 1997 (04.12.97)
(21) International Application Number: PCT/EP97/02793 (22) International Filing Date: 28 May 1997 (28.05.97) (30) Priority Data: 196 21 588.9 29 May 1996 (29.05.96) DE 196 36 917.7 11 September 1996 (11.09.96) DE (71) Applicant (for all designated States except US): HOECHST SCHERING AGREVO GMBH [DE/DE]; Miraustrasse 54, D-13509 Berlin (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): BLOCK, Martina [DE/DE]; Tiroler Strasse 13a, D-22049 Hamburg (DE). LÖRZ, Horst [DE/DE]; Ramckeweg 6a, D-22589 Hamburg (DE). LÜTTICKE, Stephanie [DE/DE]; Lange Reihe 22, D-20099 Hamburg (DE). WALTER, Lennart [DE/DE]; Am Hafen 8, D-25348 Glückstadt (DE). FROHBERG, Claus [DE/DE]; Blankenhainer Strasse 17, D-12249 Berlin (DE). KOSSMANN, Jens [DE/DE]; Golmer Fichten 9, D-14476 Golm (DE). (74) Agent: VOSSIUS & PARTNER; P.O. Box 86 07 67, D-81634 München (DE).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LU, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: NUCLEIC ACID MOLECULES ENCODING ENZYMES FROM WHEAT WHICH ARE INVOLVED IN STARCH SYNTHESIS (57) Abstract The present invention relates to nucleic acid molecules encoding enzymes which are involved in the starch synthesis in plants. These enzymes are starch synthases from wheat. The invention further relates to vectors and host cells containing said nucleic acid molecules, in particular transformed plant cells and plants regenerated from these cells, which exhibit an increased or a reduced activity of the described starch synthases.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**Nucleic acid molecules encoding enzymes from wheat which are
involved in starch synthesis**

The present invention relates to nucleic acid molecules encoding enzymes from wheat which are involved in the starch synthesis of plants. These enzymes are isotypes of the starch synthase.

The invention further relates to vectors and bacteria which contain these nucleic acid molecules as well as plant cells and plants transformed with the described nucleic acid molecules. Furthermore, methods for the production of transgenic plants are described which due to the integration of DNA molecules encoding starch synthase from wheat, synthesize starch which is modified in its properties.

With respect to its increasing significance which has recently been ascribed to vegetal substances as regenerative sources of raw materials, one of the objects of biotechnological research is to try to adapt vegetal raw materials to the demands of the processing industry. In order to enable the use of modified regenerative raw materials in as many areas as possible, it is furthermore important to obtain a large variety of substances. Apart from oils, fats and proteins, polysaccharides constitute the essential regenerative raw materials derived from plants. Apart from cellulose, starch maintains an important position

among the polysaccharides, being one of the most significant storage substances in higher plants. Among those, wheat is an interesting cultivated plant as it generates 20% of the total amount of starch produced in the European Community.

The polysaccharide starch is a polymer made up of chemically homogeneous basic components, namely the glucose molecules. However, it constitutes a highly complex mixture from various types of molecules which differ from each other in their degree of polymerization and in the degree of branching of the glucose chains. Therefore, starch is not a homogeneous raw material. One differentiates particularly between amylose-starch, a basically non-branched polymer made up of α -1,4-glycosidically branched glucose molecules, and amylopectin-starch which in turn is a complex mixture of various branched glucose chains. The branching results from additional α -1,6-glycosidic interlinkings. In wheat the synthesized starch consists of about 11-37% of amylose-starch, depending on the cultivar.

In order to enable as wide a use of starch as possible, it seems to be desirable that plants be provided which are capable of synthesizing modified starch which is particularly suitable for various uses. Breeding is one possibility to provide such plants. This, however, turns out to be very difficult in the case of wheat due to the polyploid properties of cultivated wheat (tetra- and hexaploid). Only recently scientists succeeded in producing "waxy" (not containing amylose) wheat by cross-breeding of mutants occurring in nature (Nakamura et al., Mol. Gen. Genet. 248 (1995), 253-259). Another possibility is the specific genetic modification of the starch metabolism of starch-producing plants by means of recombinant DNA techniques. However, a prerequisite therefor is to identify and to characterize the enzymes involved in the starch synthesis

and/or the starch modification as well as to isolate the respective DNA molecules encoding these enzymes.

The biochemical pathways which lead to the production of starch are basically known. The starch synthesis in plant cells takes place in the plastids. In photosynthetically active tissues these are the chloroplasts, in photosynthetically inactive, starch-storing tissues the amyloplasts.

The most important enzymes involved in starch synthesis are starch synthases as well as branching enzymes. In the case of starch synthases various isotypes are described which all catalyze a polymerization reaction by transferring a glucosyl residue of ADP-glucose to α -1,4-glucans. Branching enzymes catalyze the introduction of α -1,6 branchings into linear α -1,4-glucans.

Starch synthases may be divided up in two groups: the granule-bound starch synthases (GBSS) and the soluble starch synthases (SSS). This distinction is not always evident since some starch synthases are granule-bound as well as soluble (Denyer et al., Plant J. 4 (1993), 191-198; Mu et al., Plant J. 6 (1994), 151-159). Within these classifications, various isotypes are described for various species of plants. These isotypes differ from each other in their dependency on primer molecules (so-called "primer dependent" (type II) and "primer independent" (type I) starch synthases).

So far only in the case of the isotype GBSS I its exact function during starch synthesis has been successfully determined. Plants in which this enzyme activity has been strongly or completely reduced, synthesize starch free of amylose (a so-called "waxy" starch) (Shure et al., Cell 35 (1983), 225-233; Visser et al., Mol. Gen. Genet. 225 (1991), 289-296; WO 92/11376); therefore this enzyme has been assigned a decisive role in synthesizing amylose-starch. This phenomenon

is also observed in the cells of the green alga *Chlamydomonas reinhardtii* (Delrue et al., J. Bacteriol. 174 (1992), 3612-3620). In the case of *Chlamydomonas* it was furthermore demonstrated that GBSS I is not only involved in the synthesis of amylose but also has an influence on amylopectin synthesis. In mutants which do not show any GBSS I activity a certain fraction of the normally synthesized amylopectin, exhibiting long chain glucans, is missing.

The functions of the other isotypes of the granule-bound starch synthases, particularly GBSS II, and of the soluble starch synthases are so far not clear. It is assumed that soluble starch synthases, together with branching enzymes, are involved in the synthesis of amylopectin (see e.g. Ponstein et al., Plant Physiol. 92 (1990), 234-241) and that they play an important role in the regulation of starch synthesis rate.

In the case of wheat at least two isotypes of granule-bound starch synthase (60 kDa and 100-105 kDa) and a further isotype, which possibly represents a soluble starch synthase (Denyer et al., Planta 196 (1995), 256-265; Rahman et al., Aust. J. Plant Physiol. 22 (1995), 793-803), were identified on the protein level. The existence of several SSS-isotypes had already been proved by means of chromatographic methods (Rijven, Plant Physiol. 81 (1986), 448-453). A cDNA encoding GBSS I from wheat has already been described (Ainsworth et al., Plant Mol. Biol. 22 (1993), 67-82).

Nucleic acid sequences encoding further starch synthase-isotypes from wheat are yet unknown.

cDNA-sequences encoding other starch synthases than GBSS I have so far merely been described for pea (Dry et al., Plant J. 2 (1992), 193-202), rice (Baba et al., Plant Physiol. 103 (1993), 565-573) and potatoes (Edwards et al., Plant J. 8 (1995), 283-294).

Soluble starch synthases have been identified in several other plant species apart from wheat. Soluble starch synthases have for example been isolated in homogeneous form from pea (Denyer and Smith, *Planta* 186 (1992), 609-617) and potatoes (Edwards et al., *Plant J.* 8 (1995), 283-294). In these cases it was found that the isotype of the soluble starch synthase identified as SSS II is identical with the granule-bound starch synthase GBSS II (Denyer et al., *Plant J.* 4 (1993), 191-198; Edwards et al., *Plant J.* 8 (1995), 283-294). In the case of other plant species the existence of several SSS-isotypes was described by means of chromatographic methods, as for example in the case of barley (Tyynelä and Schulman, *Physiologia Plantarum* 89 (1993) 835-841; Kreis, *Planta* 148 (1980), 412-416). However, DNA sequences encoding these proteins have so far not been described.

In order to provide further possibilities for modifying any desired starch-storing plant, especially wheat, in such a way that they will synthesize a modified starch, respective DNA sequences encoding further isotypes of starch synthases have to be identified.

Therefore, it was the object of the present invention to provide nucleic acid molecules encoding enzymes - especially enzymes from wheat - involved in starch biosynthesis and by means of which genetically modified plants may be produced that show an elevated or reduced activity of those enzymes, thereby prompting a modification in the chemical and/or physical properties of the starch synthesized in these plants.

This object has been achieved by the provision of the embodiments described in the claims.

Therefore, in a first aspect the present invention relates to nucleic acid molecules encoding proteins from wheat with the biological activity of a soluble starch synthase, whereby such

molecules preferably encode proteins which comprise the amino acid sequence depicted under Seq ID No. 2. The invention particularly relates to nucleic acid molecules which comprise all or part of the nucleotide sequence mentioned under Seq ID No. 1, preferably molecules, which comprise the coding region indicated in Seq ID No. 1 or, as the case may be, corresponding ribonucleotide sequences.

The present invention further relates to nucleic acid molecules encoding soluble starch synthase from wheat and hybridizing to one of the above-mentioned molecules.

Nucleic acid molecules that encode soluble starch synthase from wheat and the sequence of which differs from the nucleotide sequences of the above-mentioned molecules due to the degeneracy of the genetic code are also the subject-matter of the invention.

The invention also relates to nucleic acid molecules showing a sequence which is complementary to the whole or to a part of the above-mentioned sequences.

The proteins encoded by the above-described nucleic acid molecules are soluble starch synthases derived from wheat. These proteins show certain homologous regions with the so far known soluble starch synthases from other plant species.

In another aspect the present invention relates to nucleic acid molecules encoding proteins with the biological activity of a starch synthase from wheat, whereby such molecules preferably encode proteins comprising the amino acid sequence indicated under Seq ID No. 6. The invention particularly relates to nucleic acid molecules which contain the nucleotide sequence indicated under Seq ID No. 5 or part of it, preferably molecules comprising the coding region depicted under Seq ID

No. 5 or, as the case may be, correspond ribonucleotide sequences.

The present invention further relates to nucleic acid molecules encoding starch synthase from wheat and hybridizing to one of the above mentioned molecules.

Nucleic acid molecules that encode a starch synthase from wheat and the sequence of which differs from the nucleic acid sequences of the above-described molecules due to the degeneracy of the genetic code are also the subject-matter of the invention.

The invention also relates to nucleic acid molecules showing a sequence which is complementary to the whole or to a part of the above-mentioned sequences.

The protein encoded by the above-described nucleic acid molecules is a protein with the biological activity of a starch synthase from wheat. When comparing the homology with other known sequences it was found that the highest degree of homology occurs with peas, which encode a granule-bound starch synthase. Thus, it is assumed that the described nucleic acid molecules encode a granule bound starch synthase from wheat.

The nucleic acid molecules of the invention may be DNA as well as RNA molecules. Corresponding DNA molecules are for instance genomic or cDNA molecules. The nucleic acid molecules of the invention may be isolated from natural sources or synthesized by means of known methods.

In this invention the term "hybridization" signifies hybridization under conventional hybridizing conditions, preferably under stringent conditions as described for example in Sambrook et al., Molecular Cloning, A Laboratory Manual, 2nd

Edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

Nucleic acid molecules hybridizing to the molecules according to the invention may be isolated e.g. from genomic or from cDNA libraries produced from wheat tissue.

Thereby, the identification and isolation of such nucleic acid molecules may take place by using the molecules according to the invention or parts of these molecules or, as the case may be, the reverse complement strands of these molecules, e.g. by hybridization according to standard methods (see e.g. Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, 2nd Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

As a probe for hybridization e.g. nucleic acid molecules may be used which exactly or basically contain the nucleotide sequences indicated under Seq ID No. 1 or under Seq ID No. 5 or parts thereof. The fragments used as hybridization probe may also be synthetic fragments which were produced by means of the conventional synthesizing methods and the sequence of which is basically identical with that of a nucleic acid molecule according to the invention. After identifying and isolating the genes hybridizing to the nucleic acid sequences according to the invention, the sequence has to be determined and the properties of the proteins encoded by this sequence have to be analyzed.

The molecules hybridizing to the nucleic acid molecules of the invention also comprise fragments, derivatives and allelic variants of the above-described nucleic acid molecules which encode a protein from wheat as described in the invention. Thereby, fragments are defined as parts of the nucleic acid molecules, which are long enough in order to encode one of the described proteins. This includes also parts of nucleic acid

molecules according to the invention which lack the nucleotide sequence encoding the signal peptide responsible for the translocation of the protein into the plastid. Such fragments are, for example, the nucleotide sequence encoding amino acid residues 34 to 671 as shown in Seq ID No. 2 or the nucleotide sequence encoding the amino acid residues 58 to 799 or 61 to 799 as shown in Seq ID No. 6. Furthermore, fragments which are particularly preferred in the present invention are the fragments comprising nucleotides 186 to 2239 of Seq ID No. 1 as well as fragments comprising an additional G residue at their 5'-end, and fragments comprising nucleotides 1084 to 2825 of Seq ID No. 2. In this context, the term derivatives means that the sequences of these molecules differ from the sequences of the above-mentioned nucleic acid molecules at one or more positions and that they exhibit a high degree of homology to these sequences. Hereby, homology means a sequence identity of at least 40%, in particular an identity of at least 60%, preferably of more than 80% and still more preferably a sequence identity of more than 90%. The deviations occurring when comparing with the above-described nucleic acid molecules might have been caused by deletion, substitution, insertion or recombination.

Moreover, homology means that functional and/or structural equivalence exists between the respective nucleic acid molecules or the proteins they encode. The nucleic acid molecules, which are homologous to the above-described molecules and represent derivatives of these molecules, are generally variations of these molecules, that constitute modifications which exert the same biological function. These variations may be naturally occurring variations, for example sequences derived from other organisms, or mutations, whereby these mutations may have occurred naturally or they may have

been introduced by means of a specific mutagenesis. Moreover the variations may be synthetically produced sequences. The allelic variants may be naturally occurring as well as synthetically produced variants or variants produced by recombinant DNA techniques.

The proteins encoded by the various variants of the nucleic acid molecules according to the invention exhibit certain common characteristics. Enzyme activity, molecular weight, immunologic reactivity, conformation etc. may belong to these characteristics as well as physical properties such as the mobility in gel electrophoresis, chromatographic characteristics, sedimentation coefficients, solubility, spectroscopic properties, stability, pH-optimum, temperature-optimum etc. Significant characteristics of a starch synthase are: i) their localization within the stroma of the plastids of plant cells; ii) their capability of synthesizing linear α -1,4-linked polyglucans using ADP-glucose as substrate. This activity can be determined as shown in Denyer and Smith (Planta 186 (1992), 606-617) or as described in the examples.

Nucleic acid molecules hybridizing specifically to a strand of the nucleic acid molecules of the invention are also subject-matter of the invention. These are preferably oligonucleotides with a length of at least 10, particularly of at least 15 and still more preferably with a length of at least 50 nucleotides. These nucleic acid molecules hybridize specifically to a strand of a nucleic acid molecule of the invention, i.e. they do not or only to a small extent hybridize to nucleic acid sequences encoding other proteins, particularly other starch synthases. The oligonucleotides of the invention may be used for example as primer for a PCR reaction. They may also be components of

antisense-constructs or DNA molecules encoding suitable ribozymes.

Furthermore, the invention relates to vectors, especially plasmids, cosmids, viruses, bacteriophages and other vectors common in genetic engineering, which contain the above-mentioned nucleic acid molecules of the invention.

In a preferred embodiment the nucleic acid molecules contained in the vectors are linked to regulatory elements that ensure the transcription and synthesis of a translatable RNA in procaryotic or eucaryotic cells.

The expression of the nucleic acid molecules of the invention in procaryotic cells, e.g. in *Escherichia coli*, is interesting insofar as this enables a more precise characterization of the enzymatic activities of the enzymes encoding these molecules. In particular, it is possible to characterize the product being synthesized by the respective enzymes in the absence of other enzymes which are involved in the starch synthesis of the plant cell. This makes it possible to draw conclusions about the function, which the respective protein exerts during the starch synthesis within the plant cell.

Moreover, it is possible to introduce various mutations into the nucleic acid molecules of the invention by means of conventional molecular-biological techniques (see e.g. Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY), whereby the synthesis of proteins with possibly modified biological properties is induced. By means of this it is on the one hand possible to produce deletion mutants, in which nucleic acid molecules are produced by continuing deletions at the 5'- or the 3'-end of the encoding DNA-

sequence. These nucleic acid molecules may lead to the synthesis of correspondingly shortened proteins. Such deletions at the 5'-end of the nucleotide sequence make it possible, for example, to identify amino acid sequences which are responsible for the translocation of the enzyme in the plastids (transit peptides). This allows for the specific production of enzymes which due to the removal of the respective sequences are no longer located in the plastids but within the cytosol, or which due to the addition of other signal sequences are located in other compartments.

On the other hand point mutations might also be introduced at positions where a modification of the amino acid sequence influences, for example, the enzyme activity or the regulation of the enzyme. In this way e.g. mutants with a modified K_m -value may be produced, or mutants which are no longer subject to the regulation mechanisms by allosteric regulation or covalent modification usually occurring in cells.

Furthermore, mutants may be produced exhibiting a modified substrate or product specificity such as mutants that use ADP-glucose-6-phosphate instead of ADP-glucose as substrate. Moreover, mutants with a modified activity-temperature-profile may be produced.

For the genetic manipulation in procaryotic cells the nucleic acid molecules of the invention or parts of these molecules may be integrated into plasmids which allow for a mutagenesis or a sequence modification by recombination of DNA sequences. By means of standard methods (cf. Sambrook et al., 1989, Molecular Cloning: A laboratory manual, 2nd edition, Cold Spring Harbor Laboratory Press, NY, USA) base exchanges may be carried out or natural or synthetic sequences may be added. In order to connect the DNA fragments, adapters or linkers may be attached

to the fragments. Moreover, use can be made of manipulations which offer suitable restriction sites or which remove superfluous DNA or restriction sites. Wherever use is made of inserts, deletions or substitutions, in vitro mutagenesis, "primer repair", restriction or ligation may be used. For analyzing use is usually made of a sequence analysis, a restriction analysis and further biochemico-molecularbiological methods.

In a further embodiment the invention relates to host cells, in particular procaryotic or eucaryotic cells, which have been transformed and/or genetically modified by an above-mentioned nucleic acid molecule of the invention or by a vector of the invention, as well as cells derived from cells transformed and/or genetically modified in such a way and containing a nucleic acid molecule of the invention or a vector of the invention. This is preferably a bacterial cell or a plant cell. Such cells are characterized in that the introduced nucleic acid molecule of the invention is either heterologous with respect to the transformed cell, i.e. it does not occur naturally in these cells, or is located at another place in the genome than the corresponding, naturally occurring sequence. Furthermore, the proteins encoded by the nucleic acid molecules of the invention are the subject-matter of the invention as well as methods for their production whereby a host cell of the invention is cultivated under conditions that allow for a synthesis of the protein and whereby the protein is then isolated from the cultivated cells and/or the culture medium.

Moreover, the present invention also relates to transgenic plant cells transformed with one or more nucleic acid molecule(s) of the invention. Such cells contain one or more

nucleic acid molecule(s) of the invention, whereby this/these is/are preferably linked to regulatory DNA elements, which ensure the transcription in plant cells, especially with a promoter. Such cells differ from naturally occurring plant cells in that they contain at least one nucleic acid molecule of the invention which does not naturally occur in such cells or in that such a molecule is integrated at some position in the genome of the cell at which it does not naturally occur, i.e. in a different genomic environment.

By means of methods known to the skilled person the transgenic plant cells can be regenerated to whole plants. Thus, the plants obtained by regenerating the transgenic plant cells of the invention are also the subject-matter of the present invention. A further subject-matter of the invention are plants which contain the above-described transgenic plant cells. The transgenic plants may in principle be plants of any desired species, i.e. they may be monocotyledonous as well as dicotyledonous plants. These are preferably useful plants, in particular starch-synthesizing or starch-storing plants such as cereals (rye, barley, oats, wheat etc.), rice, maize, peas, cassava or potatoes.

By making available the nucleic acid molecules of the invention it is now possible - by means of recombinant DNA techniques - to specifically interfere with the starch metabolism of plants in a way so far impossible by means of breeding. Thereby, the starch metabolism may be modified in such a way that a modified starch is synthesized which e.g. is modified, compared to the starch synthesized in wildtype plants, with respect to its physico-chemical properties, especially the amylose/amylopectin ratio, the degree of branching, the average chain length, the phosphate content, the pastification behavior, the size and/or the shape of the starch granule. There is the possibility of

increasing the yield of genetically modified plants by increasing the activity of the proteins described in the invention, e.g. by overexpressing the respective nucleic acid molecules or by making mutants available which are no longer subject to cell-specific regulation schemes and/or different temperature-dependencies with respect to their activity. The economic significance of the chance to interfere with the starch synthesis of wheat is obvious since this plant produces considerable amounts of starch.

Therefore it is possible to express the nucleic acid molecules of the invention in plant cells in order to increase the activity of the respective starch synthases or it is possible to introduce them into cells that usually do not express said enzyme. Furthermore, the nucleic acid molecules of the invention may be modified by means of methods known to the skilled person, in order to produce starch synthases according to the invention which are no longer subject to the cell-specific regulation mechanisms or show modified temperature-dependencies or substrate resp. product specificities.

In expressing the nucleic acid molecules of the invention in plants the synthesized proteins may in principle be located in any desired compartment within the plant cell. In order to locate it within a specific compartment, the sequence ensuring the localization in the plastids must be deleted and the remaining coding regions optionally have to be linked to DNA sequences which ensure localization in the respective compartment. Such sequences are known (see e.g. Braun et al., EMBO J. 11 (1992), 3219-3227; Wolter et al., Proc. Natl. Acad. Sci. USA 85 (1988), 846-850; Sonnewald et al., Plant J. 1 (1991), 95-106).

The invention also relates to propagation material of the plants of the invention, e.g. fruits, seeds, tubers, root-stocks, seedlings, cuttings, calli, cell cultures etc.

The starch derived from transgenic plant cells, plants as well as the propagation material according to the invention is also the subject-matter of the present invention.

Due to the expression or, as the case may be, additional expression of at least one of the nucleic acid molecules of the invention, the transgenic plant cells and plants described in the invention synthesize a starch which compared to starch synthesized in wildtype plants is modified for example in its physico-chemical properties, in particular in the amylose/amylopectin ratio, the degree of branching, the average chain-length, the phosphate-content, the pastification behavior, the size and/or the shape of the starch granule. Compared with wildtype-starch, such starch may be modified in particular with respect to its viscosity and/or the gel formation properties of the glues of this starch.

Transgenic plant cells, in which the activity of at least one protein according to the invention is reduced when compared to non-transformed plants, are a further subject-matter of the invention.

By means of the nucleic acid molecules of the invention it is possible to produce plant cells and plants in which the activity of at least one protein of the invention is reduced. This also leads to the synthesis of a starch with modified chemical and/or physical properties when compared to the starch from wildtype plant cells.

The production of plant cells with a reduced activity of at least one protein of the invention may, for example, be achieved by the expression of at least one corresponding

antisense-RNA, of at least one sense-RNA or achieving a cosuppression effect or the expression of at least one correspondingly constructed ribozyme, which specifically cleaves transcripts encoding one of the proteins of the invention, using the nucleic acid molecules of the invention.

In order to express an antisense-RNA, on the one hand DNA molecules can be used which comprise the complete sequence encoding a protein of the invention, including possibly existing flanking sequences as well as DNA molecules, which only comprise parts of the encoding sequence whereby these parts have to be long enough in order to prompt an antisense-effect within the cells. Basically, sequences with a minimum length of 15 bp, preferably with a length of 100-500 bp and for an efficient antisense-inhibition, in particular sequences with a length of more than 500 bp may be used. Generally DNA-molecules are used which are shorter than 5000 bp, preferably sequences with a length of less than 2500 bp.

Use may also be made of DNA sequences which are highly homologous, but not completely identical to the sequences of the DNA molecules of the invention. The minimal homology should be more than about 65%. Preferably, use should be made of sequences with homologies between 95 and 100%.

The method for reducing the activity of enzymes of the invention in plant cells by means of a cosuppression effect is known to the skilled person and has been described, for example, in Jorgensen (Trends Biotechnol. 8 (1990), 340-344), Niebel et al. (Curr. Top. Microbiol. Immunol. 197 (1995), 91-103), Flavell et al. (Curr. Top. Microbiol. Immunol. 197 (1995), 43-46), Palaqui and Vaucheret (Plant. Mol. Biol. 29 (1995), 149-159), Vaucheret et al. (Mol. Gen. Genet. 248 (1995), 311-317), de Borne et al. (Mol. Gen. Genet. 243 (1994), 613-621) and in other sources.

The expression of corresponding ribozymes in order to reduce the activity of certain enzymes in cells is also known to the person skilled in the art and described, for example, in EP-B1 0 321 201. The expression of ribozymes in plant cells was described e.g. by Feyter et al. (Mol. Gen. Genet. 250 (1996), 329-338).

Moreover, plants containing the above-described transgenic plant cells of the invention are also the subject-matter of the present invention. These may be regenerated from the plant cells of the invention to whole plants by means of methods known to the skilled person. These plants are preferably those already mentioned above, in particular useful plants, especially starch-synthesizing or, as the case may be, starch-storing plants. Hereby, wheat is particularly preferred.

The invention also relates to propagation material of the plants of the invention, in particular to fruits, seeds, tubers, rootstocks, seedlings, cuttings, calli, cell cultures etc.

Moreover, starch derived from the above-mentioned transgenic plant cells, plants as well as the propagation material is the subject-matter of the invention.

Due to the reduction of the activity of at least one of the proteins of the invention, the transgenic plant cells and plants of the invention synthesize a starch which is modified, compared to the starch synthesized in wildtype plants, in its physico-chemical properties, in particular in the amylose/amylopectin ratio, the degree of branching, the average chain-length, the phosphate-content, the pastification behavior, the size and/or the shape of the starch granule. This

starch may for example exhibit modified viscosities and/or gel formation properties of its gels when compared to starch derived from wildtype plants.

The starches of the invention may be modified according to techniques known to the skilled person; in unmodified as well as in modified form they are suitable for the use in foodstuffs or for the use in non-foodstuffs.

Basically, the possibilities of uses of the starch can be subdivided into two major fields. One field comprises the hydrolysis products of starch, essentially glucose and glucans components obtained by enzymatic or chemical processes. They can be used as starting material for further chemical modifications and processes, such as fermentation. In this context, it might be of importance that the hydrolysis process can be carried out simply and inexpensively. Currently, it is carried out substantially enzymatically using amyloglucosidase. It is thinkable that costs might be reduced by using lower amounts of enzymes for hydrolysis due to changes in the starch structure, e.g. increasing the surface of the grain, improved digestibility due to less branching or a steric structure, which limits the accessibility for the used enzymes.

The other field in which the starch is used because of its polymer structure as so-called native starch, can be subdivided into two further areas:

1. Use in foodstuffs

Starch is a classic additive for various foodstuffs, in which it essentially serves the purpose of binding aqueous additives and/or causes an increased viscosity or an increased gel formation. Important characteristic properties are flowing and sorption behavior, swelling and

pastification temperature, viscosity and thickening performance, solubility of the starch, transparency and paste structure, heat, shear and acid resistance, tendency to retrogradation, capability of film formation, resistance to freezing/thawing, digestibility as well as the capability of complex formation with e.g. inorganic or organic ions.

A preferred area of application of native starch is the field of bakery-goods and pasta.

2. Use in non-foodstuffs

The other major field of application is the use of starch as an adjuvant in various production processes or as an additive in technical products. The major fields of application for the use of starch as an adjuvant are, first of all, the paper and cardboard industry. In this field, the starch is mainly used for retention (holding back solids), for sizing filler and fine particles, as solidifying substance and for dehydration. In addition, the advantageous properties of starch with regard to stiffness, hardness, sound, grip, gloss, smoothness, tear strength as well as the surfaces are utilized.

2.1 Paper and cardboard industry

Within the paper production process, a differentiation can be made between four fields of application, namely surface, coating, mass and spraying.

The requirements on starch with regard to surface treatment are essentially a high degree of brightness, corresponding viscosity, high viscosity stability, good film formation as well as low formation of dust. When used in coating the solid content, a corresponding viscosity, a

high capability to bind as well as a high pigment affinity play an important role. As an additive to the mass rapid, uniform, loss-free dispersion, high mechanical stability and complete retention in the paper pulp are of importance. When using the starch in spraying, corresponding content of solids, high viscosity as well as high capability to bind are also significant.

2.2 Adhesive industry

A major field of application is, for instance, in the adhesive industry, where the fields of application are subdivided into four areas: the use as pure starch glue, the use in starch glues prepared with special chemicals, the use of starch as an additive to synthetic resins and polymer dispersions as well as the use of starches as extenders for synthetic adhesives. 90% of all starch-based adhesives are used in the production of corrugated board, paper sacks and bags, composite materials for paper and aluminum, boxes and wetting glue for envelopes, stamps, etc.

2.3 Textile and textile care industry

Another possible use as adjuvant and additive is in the production of textiles and textile care products. Within the textile industry, a differentiation can be made between the following four fields of application: the use of starch as a sizing agent, i.e. as an adjuvant for smoothing and strengthening the burring behavior for the protection against tensile forces active in weaving as well as for the increase of wear resistance during weaving, as an agent for textile improvement mainly after quality-deteriorating pretreatments, such as bleaching,

dying, etc., as thickener in the production of dye pastes for the prevention of dye diffusion and as an additive for warping agents for sewing yarns.

2.4 Building industry

The fourth area of application of starch is its use as an additive in building materials. One example is the production of gypsum plaster boards, in which the starch mixed in the thin plaster pastifies with the water, diffuses at the surface of the gypsum board and thus binds the cardboard to the board. Other fields of application are admixing it to plaster and mineral fibers. In ready-mixed concrete, starch may be used for the deceleration of the sizing process.

2.5 Ground stabilization

Furthermore, the starch is advantageous for the production of means for ground stabilization used for the temporary protection of ground particles against water in artificial earth shifting. According to state-of-the-art knowledge, combination products consisting of starch and polymer emulsions can be considered to have the same erosion- and incrustation-reducing effect as the products used so far; however, they are considerably less expensive.

2.6 Use of starch in plant protectives and fertilizers

Another field of application is the use of starch in plant protectives for the modification of the specific properties of these preparations. For instance, starches are used for improving the wetting of plant protectives and fertilizers, for the dosed release of the active ingredients, for the conversion of liquid, volatile and/or

odorous active ingredients into microcrystalline, stable, deformable substances, for mixing incompatible compositions and for the prolongation of the duration of the effect due to a reduced disintegration.

2.7 Drugs, medicine and cosmetics industry

Starch may also be used in the fields of drugs, medicine and in the cosmetics industry. In the pharmaceutical industry, the starch may be used as a binder for tablets or for the dilution of the binder in capsules. Furthermore, starch is suitable as disintegrant for tablets since, upon swallowing, it absorbs fluid and after a short time it swells so much that the active ingredient is released. For qualitative reasons, medicinal powders and dusting powders are further fields of application. In the field of cosmetics, the starch may for example be used as a carrier of powder additives, such as scents and salicylic acid. A relatively extensive field of application for the starch is toothpaste.

2.8 Starch as an additive in coal and briquettes

The use of starch as an additive in coal and briquettes is also thinkable. By adding starch, coal can be quantitatively agglomerated and/or briquetted in high quality, thus preventing premature disintegration of the briquettes. Barbecue coal contains between 4 and 6% added starch, calorated coal between 0.1 and 0.5%. Furthermore, the starch is suitable as a binding agent since adding it to coal and briquette can considerably reduce the emission of toxic substances.

2.9 Processing of ore and coal slurry

Furthermore, the starch may be used as a flocculant in the processing of ore and coal slurry.

2.10 Starch as an additive in casting

Another field of application is the use as an additive to process materials in casting. For various casting processes cores produced from sands mixed with binding agents are needed. Nowadays, the most commonly used binding agent is bentonite mixed with modified starches, mostly swelling starches.

The purpose of adding starch is increased flow resistance as well as improved binding strength. Moreover, swelling starches may fulfill more prerequisites for the production process, such as dispersability in cold water, rehydratisability, good mixability in sand and high capability of binding water.

2.11 Use of starch in rubber industry

In the rubber industry starch may be used for improving the technical and optical quality. Reasons for this are improved surface gloss, grip and appearance. For this purpose, the starch is dispersed on the sticky rubberized surfaces of rubber substances before the cold vulcanization. It may also be used for improving the printability of rubber.

2.12 Production of leather substitutes

Another field of application for the modified starch is the production of leather substitutes.

2.13 Starch in synthetic polymers

In the plastics market the following fields of application are emerging: the integration of products derived from starch into the processing process (starch is only a filler, there is no direct bond between synthetic polymer and starch) or, alternatively, the integration of products derived from starch into the production of polymers (starch and polymer form a stable bond).

The use of the starch as a pure filler cannot compete with other substances such as talcum. This situation is different when the specific starch properties become effective and the property profile of the end products is thus clearly changed. One example is the use of starch products in the processing of thermoplastic materials, such as polyethylene. Thereby, starch and the synthetic polymer are combined in a ratio of 1 : 1 by means of coexpression to form a 'master batch', from which various products are produced by means of common techniques using granulated polyethylene. The integration of starch in polyethylene films may cause an increased substance permeability in hollow bodies, improved water vapor permeability, improved antistatic behavior, improved anti-block behavior as well as improved printability with aqueous dyes. Another possibility is the use of the starch in polyurethane foams. Due to the adaptation of starch derivatives as well as due to the optimization of processing techniques, it is possible to specifically control the reaction between synthetic

polymers and the starch's hydroxy groups. The results are polyurethane films having the following property profiles due to the use of starch: a reduced coefficient of thermal expansion, decreased shrinking behavior, improved pressure/tension behavior, increased water vapor permeability without a change in water acceptance, reduced flammability and cracking density, no drop off of combustible parts, no halides and reduced aging. Disadvantages that presently still exist are reduced pressure and impact strength.

Product development of film is not the only option. Also solid plastics products, such as pots, plates and bowls can be produced by means of a starch content of more than 50%. Furthermore, the starch/polymer mixtures offer the advantage that they are much easier biodegradable.

Furthermore, due to their extreme capability to bind water, starch graft polymers have gained utmost importance. These are products having a backbone of starch and a side lattice of a synthetic monomer grafted on according to the principle of radical chain mechanism. The starch graft polymers available nowadays are characterized by an improved binding and retaining capability of up to 1000 g water per g starch at a high viscosity. These super absorbers are used mainly in the hygiene field, e.g. in products such as diapers and sheets, as well as in the agricultural sector, e.g. in seed pellets.

What is decisive for the use of the new starch modified by recombinant DNA techniques are, on the one hand, structure, water content, protein content, lipid content, fiber content, ashes/phosphate content, amylose/amylopectin ratio, distribution of the relative molar mass, degree of branching, granule size and shape as well as crystallization, and on the other hand, the properties resulting in the following features:

flow and sorption behavior, pastification temperature, viscosity, thickening performance, solubility, paste structure, transparency, heat, shear and acid resistance, tendency to retrogradation, capability of gel formation, resistance to freezing/thawing, capability of complex formation, iodine binding, film formation, adhesive strength, enzyme stability, digestibility and reactivity.

The production of modified starch by genetically operating with a transgenic plant may modify the properties of the starch obtained from the plant in such a way as to render further modifications by means of chemical or physical methods superfluous. On the other hand, the starches modified by means of recombinant DNA techniques might be subjected to further chemical modification, which will result in further improvement of the quality for certain of the above-described fields of application. These chemical modifications are principally known to the person skilled in the art. These are particularly modifications by means of

- heat treatment
- acid treatment
- oxidation and
- esterification

leading to the formation of phosphate, nitrate, sulfate, xanthate, acetate and citrate starches. Other organic acids may also be used for the esterification:

- formation of starch ethers

starch alkyl ether, O-allyl ether, hydroxylalkyl ether, O-carboxymethyl ether, N-containing starch ethers, P-containing starch ethers and S-containing starch ethers.

- formation of branched starches

- formation of starch graft polymers.

The starches of the invention are preferably used in the production of packaging and disposable material.

In order to express the nucleic acid molecules of the invention in sense- or antisense-orientation in plant cells, these are linked to regulatory DNA elements which ensure the transcription in plant cells. Such regulatory DNA elements are particularly promoters. Basically any promoter which is active in plant cells may be used for the expression.

The promoter may be selected in such a way that the expression takes place constitutively or in a certain tissue, at a certain point of time of the plant development or at a point of time determined by external circumstances. With respect to the plant the promoter may be homologous or heterologous. Suitable promoters for a constitutive expression are, e.g. the 35S RNA promoter of the Cauliflower Mosaic Virus and the ubiquitin promoter from maize. For a tuber-specific expression in potatoes the patatin gene promoter B33 (Rocha-Sosa et al., EMBO J. 8 (1989), 23-29) can be used. An example for a promoter which ensures expression only in photosynthetically active tissues is the ST-LS1 promoter (Stockhaus et al., Proc. Natl. Acad. Sci. USA 84 (1987), 7943-7947; Stockhaus et al., EMBO J.

8 (1989), 2445-2451). For an endosperm-specific expression the HMG promoter from wheat, the USP promoter, the phaseolin promoter or promoters from zein genes from maize are suitable. Furthermore, a termination sequence may exist which serves to correctly end the transcription and to add a poly-A-tail to the transcript which is believed to stabilize the transcripts. Such elements are described in the literature (cf. Gielen et al., EMBO J. 8 (1989), 23-29) and can be exchanged as desired.

The present invention provides nucleic acid molecules encoding two distinctive types of starch synthase from wheat. This allows for the identification of the function of these isotypes in the starch biosynthesis as well as for the production of genetically modified plants in which the activity of at least one of these enzymes is modified. This enables the synthesis of starch with a modified structure and therefore with modified physico-chemical properties in the plants manipulated in such a way.

The nucleic acid molecules of the invention may also be used in order to produce plants in which the activity of at least one of the starch synthases of the invention is elevated or reduced and in which at the same time the activities of other enzymes involved in the starch biosynthesis are modified. Thereby, all kinds of combinations and permutations are thinkable. By modifying the activity of one or more isotypes of the starch synthases in plants, a synthesis of a starch modified in its structure is brought about. By increasing the activity of one or more isotypes of the starch synthases in the cells of the starch-storing tissue of transformed plants such as in the endosperm of maize or wheat or in the potato tuber, increased yields may be the result. For example, nucleic acid molecules encoding a protein of the invention, or corresponding antisense-constructs may be integrated into plant cells in

which the synthesis of endogenous GBSS I-, SSS- or GBSS II-proteins is already inhibited due to an antisense-effect or a mutation, or in which the synthesis of the branching enzyme is inhibited (as described e.g. in Nakamura et al. (loc. cit.)).

If the inhibition of the synthesis of several starch synthases in transformed plants is to be achieved, DNA molecules can be used for transformation, which at the same time contain several regions in antisense-orientation controlled by a suitable promoter and encoding the corresponding starch synthases. Hereby, each sequence may be controlled by its own promoter or else the sequences may be transcribed as a fusion of a common promoter. The last alternative will generally be preferred as in this case the synthesis of the respective proteins should be inhibited to approximately the same extent.

Furthermore it is possible to construct DNA molecules which, apart from DNA sequences that encode starch synthases, contain further DNA sequences encoding other proteins involved in starch synthesis or modification. Hereby, the sequences may again be connected up in series and be transcribed by a common promoter. For the length of the individual coding regions used in such a construct the above-mentioned facts concerning the production of antisense-construct are also true. There is no upper limit for the number of antisense fragments transcribed from a promoter in such a DNA molecule. The resulting transcript, however, should not be longer than 10 kb, preferably 5 kb.

Coding regions which are located in antisense-orientation behind a suitable promoter in such DNA molecules in combination with other coding regions, may be derived from DNA sequences encoding the following proteins: granule-bound starch synthases (GBSS I and II), other soluble starch synthases, branching

enzymes, debranching enzymes, disproportionating enzymes and starch phosphorylases. This enumeration merely serves as an example. The use of other DNA sequences within the framework of such a combination is also thinkable.

By means of such constructs it is possible to inhibit the synthesis of several enzymes at the same time within the plant cells transformed with these molecules.

Furthermore, the constructs may be integrated into classical mutants which are defective for one or more genes of the starch biosynthesis. These defects may be related to the following proteins: granule-bound (GBSS I and II) and soluble starch synthases (SSS I and II), branching enzymes (BE I and II), debranching enzymes (R-enzymes), disproportionizing enzymes and starch phosphorylases. This enumeration merely serves as an example.

By means of such strategy it is furthermore possible to inhibit the synthesis of several enzymes at the same time within the plant cells transformed with these nucleic acid molecules.

In order to prepare the integration of foreign genes into higher plants a high number of cloning vectors are at disposal, containing a replication signal for E.coli and a marker gene for the selection of transformed bacterial cells. Examples for such vectors are pBR322, pUC series, M13mp series, pACYC184 etc. The desired sequence may be integrated into the vector at a suitable restriction site. The obtained plasmid is used for the transformation of E.coli cells. Transformed E.coli cells are cultivated in a suitable medium and subsequently harvested and lysed. The plasmid is recovered. As an analyzing method for the characterization of the obtained plasmid DNA use is generally made of restriction analysis, gel electrophoresis and other biochemico-molecularbiological methods. After each manipulation the plasmid DNA may be cleaved and the obtained

DNA fragments may be linked to other DNA sequences. Each plasmid DNA may be cloned into the same or in other plasmids.

In order to integrate DNA into plant host cells a wide range of techniques are at disposal. These techniques comprise the transformation of plant cells with T-DNA by using *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* as transformation medium, the fusion of protoplasts, the injection and the electroporation of DNA, the integration of DNA by means of the biolistic method as well as further possibilities.

In the case of injection, the biolistic method and electroporation of DNA into plant cells, there are no special demands made to the plasmids used. Simple plasmids such as pUC derivatives may be used. However, in case that whole plants are to be regenerated from cells transformed in such a way, a selectable marker gene should be present.

Depending on the method of integrating desired genes into the plant cell, further DNA sequences may be necessary. If the Ti- or Ri-plasmid is used e.g. for the transformation of the plant cell, usually at least the right border, more frequently, however, the right and left border of the Ti- and Ri-plasmid T-DNA should be connected to the foreign gene to be integrated as a flanking region.

If *Agrobacteria* are used for the transformation, the DNA which is to be integrated should be cloned into special plasmids, namely either into an intermediate vector or into a binary vector. Due to sequences homologous to the sequences within the T-DNA, the intermediate vectors may be integrated into the Ti- or Ri-plasmid of the *Agrobacterium* due to homologous recombination. This also contains the vir-region necessary for the transfer of the T-DNA. Intermediate vectors cannot replicate in *Agrobacteria*. By means of a helper plasmid the intermediate vector may be transferred to *Agrobacterium*

tumefaciens (conjugation). Binary vectors y replicate in E.coli as well as in Agrobacteria. They contain a selectable marker gene as well as a linker or polylinker which is framed by the right and the left T-DNA border region. They may be transformed directly into the Agrobacteria (Holsters et al. Mol. Gen. Genet. 163 (1978), 181-187). The Agrobacterium acting as host cell should contain a plasmid carrying a vir-region. The vir-region is usually necessary for the transfer of the T-DNA into the plant cell. Additional T-DNA may be present. The Agrobacterium transformed in such a way is used for the transformation of plant cells.

The use of T-DNA for the transformation of plant cells was investigated intensely and described sufficiently in EP 120 516; Hoekema, In: The Binary Plant Vector System Offsetdrukkerij Kanters B.V., Alblisserdam (1985), Chapter V; Fraley et al., Crit. Rev. Plant. Sci., 4, 1-46 and An et al. EMBO J. 4 (1985), 277-287.

For transferring the DNA into the plant cells, plant explants may suitably be co-cultivated with Agrobacterium tumefaciens or Agrobacterium rhizogenes. From the infected plant material (e.g. pieces of leaves, stem segments, roots, but also protoplasts or suspension-cultivated plant cells) whole plants may then be regenerated in a suitable medium which may contain antibiotics or biozides for the selection of transformed cells. The plants obtained in such a way may then be examined as to whether the integrated DNA is present or not. Other possibilities in order to integrate foreign DNA by using the biolistic method or by transforming protoplasts are known to the skilled person (cf. e.g. Willmitzer, L., 1993 Transgenic plants. In: Biotechnology, A Multi-Volume Comprehensive Treatise (H.J. Rehm, G. Reed, A. Pühler, P. Stadler, editors), Vol. 2, 627-659, VCH Weinheim-New York-Basel-Cambridge).

Alternative Systems for the transformation of monocotyledonous plants are the transformation by means of a biolistic approach, the electrically or chemically induced DNA integration in protoplasts, the electroporation of partially permeabilized cells, the macro-injection of DNA into inflorescences, the micro-injection of DNA into microspores and pro-embryos, the DNA integration by sprouting pollen and the DNA integration in embryos by swelling (review given in: Potrykus, *Physiol. Plant* (1990), 269-273).

Whereas the transformation of dicotyledonous plants by Ti-plasmid-vector systems by means of *Agrobacterium tumefaciens* is a well-established method, more recent studies indicate that the transformation with vectors based on *Agrobacterium* can also be used in the case of monocotyledonous plants (Chan et al., *Plant Mol. Biol.* 22 (1993), 491-506; Hiei et al., *Plant J.* 6 (1994), 271-282; Bytebier et al., *Proc. Natl. Acad. Sci. USA* 84 (1987), 5345-5349; Raineri et al., *Bio/Technology* 8 (1990), 33-38; Gould et al., *Plant Physiol.* 95 (1991), 426-434; Mooney et al., *Plant, Cell Tiss. & Org. Cult.* 25 (1991), 209-218; Li et al., *Plant Mol. Biol.* 20 (1992), 1037-1048).

Three of the above-mentioned transformation systems have in the past been established for various types of cereals: electroporation of plant tissue, transformation of protoplasts and the DNA-transfer by particle-bombardment in regenerative tissue and cells (review given in: Jähne et al., *Euphytica* 85 (1995), 35-44).

In the corresponding literature the transformation of wheat is described in various ways (reviewed in Maheshwari et al., *Critical Reviews in Plant Science* 14 (2) (1995), 149-178).

Hess et al. (*Plant Sci.* 72 (1990), 233) used macroinjection in order to bring pollen and *Agrobacteria* close to each other. The mobilization of the plasmid containing the nptII gene as

selectable marker was proved by means of Southern blot analysis and the NPTII test. The transformants constituted a normal phenotype and were fertile. The kanamycin-resistance could be proved in two successive generations.

The first transgenic, fertile wheat plant that could be regenerated after its bombardment with microprojectile-bound DNA was described in Vasil et al. (Bio/Technology 10 (1992), 667-674). The target tissue for the bombardment was an embryogenic callus culture (type C callus). The bar gene, encoding a phosphinotricine phosphotransferase and therefore conveying a resistance against the herbicide phosphinotricine, was used as selectable marker gene.

A further system was described by Weeks et al. (Plant Physiol. 102 (1993), 1077-1084) as well as Becker et al. (Plant J. 5(2) (1994), 299-307). Here the scutellum of immature embryos was used as target tissue for the DNA transformation. In an introductory in vitro phase the scutellum had been made to induce somatic embryos. The efficiency of the transformation is considerably higher in the system developed by Becker et al. (loc. cit.), with 1 transgenic plant per 83 embryos of the 'Florida' kind, than in the system established by Weeks et al., with 1 to 2 transgenic plants per 1000 embryos of the 'Bobwhite' kind.

The system developed by Becker et al. (loc. cit.) constitutes a basis for the transformation experiments described in the examples.

Once the introduced DNA has been integrated in the genome of the plant cell, it usually continues to be stable there and also remains within the descendants of the originally transformed cell. It usually contains a selectable marker which confers resistance against a biozide such as phosphinotricine or against an antibiotic such as kanamycin, G 418, bleomycin or

hygromycin etc. to the transformed plant cells. The individually selected marker should therefore allow for a selection of transformed cells to cells lacking the integrated DNA.

The transformed cells grow in the usual way within the plants (see also McCormick et al., Plant Cell Reports 5 (1986), 81-84). The resulting plants can be cultivated in the usual way and cross-bred with plants having the same transformed genetic heritage or another genetic heritage. The resulting hybrid individuals have the corresponding phenotypic properties. The plant cells produce seeds.

Two or more generations should be grown in order to ensure whether the phenotypic feature is kept stably and whether it is transferred. Furthermore, seeds should be harvested in order to ensure that the corresponding phenotype or other properties will remain.

In the examples use is made of the following methods:

1. Cloning

For cloning in E.coli the vector pBluescript II SK (Stratagene) was used.

2. Bacterial strains

For the Bluescript vector and for the antisense-constructs use was made of the E.coli strain DH5 α (Bethesda Research Laboratories, Gaithersburgh, USA). For the in vivo excision the E.coli strain XL1-Blue was used.

3. Transformation of immature wheat embryos

Media used

MS: 100 ml/l macrosalts
 1 ml/l microsalts
 2 ml/l Fe/NaEDTA
 30 g/l sucrose

(D. Becker and H. Lörz, Plant Tissue Culture Manual
(1996), B12: 1-20)

#30: MS + 2.4-D (2 mg/l)

#31: MS + 2,4-D (2 mg/l) + phosphinotricine (PPT,
active component of the
herbicide BASTA® (2 mg/l)

#32: MS + 2.4-D (0,1 mg/l) + PPT (2 mg/l)

#39: MS + 2.4-D (2 mg/ml)
+ each 0.5 M mannitol/sorbitol

The indicated media were adjusted to a pH value of 5.6 with KOH and reinforced with 0.3 % of Gelrite.

The method for transforming immature embryos from wheat was developed and optimized by Becker and Lörz (D. Becker and H. Lörz, Plant Tissue Culture Manual (1996), B12: 1-20).

In the experiments described in the following the protocol laid down by Becker and Lörz (loc. cit.) was observed.

For the transformation ears with caryopses in the developing stage of 12 to 14 days are harvested after anthesis and subjected to surface sterilization. The isolated scutella are plated with the embryo axis facing the medium on the induction medium #30.

After 2-4 days of preculturing (26°C, dark) the explantates are transferred to medium #39 for osmotic preculturing (2-4 h, 26°C, dark).

For biolistic transformation 29µg of gold particles onto which 5µg or 73 ng of the target-DNA have been precipitated are used for each shot. As the experiments carried out are co-transformations, the target-DNA is added to the precipitation mixture in a proportion of 1:1, consisting of the target gene and a resistance marker gene (bar-gene).

4. DIG-labelling of the DNA fragments

The labelling of DNA fragments used as screening probes was achieved by a specific PCR by incorporating a DIG-labelled dUTP (Boehringer Mannheim, Germany).

Example 1

Identification, isolation and characterization of a cDNA encoding soluble starch synthase from wheat (*Triticum aestivum* L. cv Florida)

The synthesis of cDNA resulted from poly(A)⁺-RNA of approximately 21 day-old wheat caryopses. All experiments mentioned in the following were carried out according to the protocol of the manufacturer (ZAP-cDNA Synthesis Kit and ZAP-cDNA Gigapack II Gold Cloning Kit, Stratagene GmbH, Heidelberg).

After determining the titers of the cDNA library a primary titer of 1.25×10^6 pfu/ml could be found. The screening was carried out by means of a DIG-labelled DNA fragment. Hereby, a DIG-labelled PCR fragment encoding a subfragment from the soluble starch synthase from rice (Baba et al., loc. cit.) was used as a probe. The primers used for the PCR had the sequence

R₁: ACA GGA TCC TGT GCT ATG CGG CGT GTG AAG (Seq. ID No. 3)

R₂: TTG GGA TCC GCA ATG CCC ACA GCA TTT TTT TC (Seq. ID No. 4)

For screening approximately 5×10^4 pfu per plate (15 cm in diameter) were plated. Positive clones were singled out. By means of in vivo excision singled-out clones were obtained as pBluescript SK (-) phagemides.

After analyzing the clones by means of mini preparations and after restriction of the plasmid-DNA the TaSSS clone was further processed.

Example 2

Sequence analysis of the cDNA insert of the pTaSSS plasmid

The plasmid DNA of the clone TaSSS was isolated and the sequence of the cDNA insert was determined by means of the dideoxynucleotide-method (Sanger et al., Proc. Natl. Acad. Sci. USA 74 (1977), 5463-5467).

First, a partial sequence was determined comprising nucleotides 186 to 2239 as depicted in Seq ID No. 1 which contained an additional G residue at its 5'-end. The insert of the clone TaSSS has a length of 2239 bp and constitutes a nearly full-length cDNA. The nucleotide sequence is indicated under Seq ID No. 1. The corresponding amino acid sequence is indicated under Seq ID No. 2. A putative signal peptide cleavage site is located between amino acid residues 33 and 34 indicated in Seq ID No. 1.

A sequence analysis and a comparison with already published sequences showed that the sequence shown under Seq ID No. 1 is new and comprises a nearly full-length coding region which exhibits homologies to soluble starch synthases from other organisms. By means of the partial cDNA sequence of TaSSS it is possible for the person skilled in the field of molecular biology to isolate the missing region at the 5'-region and thereby to obtain a complete cDNA clone. In order to do so the 5'-region of the clone TaSSS may be used as probe for screening for the whole cDNA and a complete clone may be isolated using standard methods by means of hybridization. On the other hand the missing 5'-end may be obtained by using a 5'-Race-method (e.g. of Boehringer Mannheim or other manufacturers).

Example 3

Producing the plant transformation vector pTaSSS-as

In order to express an antisense-RNA to the isolated cDNA from wheat a plant transformation vector was designed on the basis of pUC19 as base plasmid in which the cDNA insert of the plasmid pTaSSS is linked to a DNA fragment in antisense-

orientation, whereby the expression is regulated by the ubiquitin-promoter. This promoter consists of the first untranslated exon and the first intron of the ubiquitin1 gene from maize (Christensen A.H. et al., Plant Molecular Biology 18 (1992), 675-689).

Parts of the polylinker and the NOS-terminator are obtained from the plasmid pAct1.cas (CAMBIA, TG 0063; Cambia, GPO Box 3200, Canberra ACT 2601, Australia). Vector constructs with this terminator and constructs based on pAct1.cas are described in McElroy et al. (Molecular Breeding 1 (1995), 27-37). For the transformation of wheat pTaSSS was used as described above.

Example 4

Identification, isolation and characterization of another cDNA encoding starch synthase from wheat (*Triticum aestivum* L., cv Florida)

In a sequence comparison of the so far known sequences encoding soluble and granule-bound starch synthases from plants, it was obvious that there are three strongly conserved regions in between the various proteins.

In order to isolate soluble starch synthases from wheat, these three regions were selected in order to generate polyclonal peptide antibodies. Therefore three synthetic polypeptides with the following amino acid sequences were produced:

Peptide 1:	NH ₂ -PWSKTGGLGDVC-COOH	(SEQ ID NO: 7)
Peptide 2:	NH ₂ -PSRFEPGLNQLY-COOH	(SEQ ID NO: 8)
Peptide 3:	NH ₂ -GTGGLRDTVENC-COOH	(SEQ ID NO: 9)

These peptides were coupled to a KLH carrier (keyhole limpet homocyanin) and subsequently used for the production of polyclonal antibodies in rabbits (Eurogentec, Seraing, Belgium).

The resulting antibodies were designated as follows:

anti-SS1 polyclonal antibody against peptide 1

anti-SS2 polyclonal antibody against peptide 2

anti-SS3 polyclonal antibody against peptide 3.

The antibodies were subsequently used in order to screen a cDNA library from wheat caryopses for sequences encoding starch synthases from wheat. For this purpose a cDNA expression library produced as described in example 1 was used. For the analysis of the phage plaques, these were transferred to nitrocellulose filters which had previously been incubated in a 10 mM IPTG solution for 30-60 minutes and subsequently been dried on Whatman paper. The transfer took 3 h at 37°C. Afterwards the filter were incubated in a blocking solution for 30 min at room temperature and washed twice in TBST-puffer for 5-10 min. The filters were shaken with the polyclonal antibodies in a suitable dilution for 1 h at room temperature or for 16 h at 4°C. The identification of plaques expressing a protein which had been recognized by one of the antibodies was carried out by means of the Immun-Blot Assay Kit; Goat Anti-Rabbit IgG (Biorad) according to manufacturer's specification.

Phage clones of the cDNA library expressing a protein which had been recognized by one of the antibodies were further purified by using standard methods. By means of the in vivo excision-method (Stratagene) E.coli-clones were produced from positive phage clones, which contained a doublestranded pBluescript II SK plasmid with the corresponding cDNA insert between the EcoRI and the XhoI site of the polylinker. After checking the size

and the restriction pattern of the insert suitable clone, TaSS1 was subjected to a sequence analysis.

Example 5

Sequence analysis of the cDNA inserts of the pTaSS1 plasmid

The plasmid-DNA was isolated from the pTaSS1 clone and the sequence of the cDNA insert was determined by means of standard methods using the dideoxynucleotide method (Sanger et al., Proc. Natl. Acad. Sci. USA 74 (1977), 5463-5467).

First, a partial sequence was determined comprising the nucleotides 1084 to 2825 as depicted in Seq ID No. 5.

The insert of the pTaSS1 clone has a length of 2825 bp and constitutes a complete cDNA. The nucleotide sequence is indicated under Seq ID No. 5. The corresponding amino acid sequence is indicated under Seq ID No. 6.

A sequence analysis and a comparison with already published sequences showed that the sequence indicated under Seq ID No. 5 is new and comprises a coding region exhibiting homologies to starch synthases from other organisms. It is assumed that this cDNA encodes a protein having the biological activity of a granule bound starch synthase.

Furthermore, due to homologies with known consensus sequences for signal peptide cleavage sites it had been found that the putative signal transit peptide is cleaved of between positions 57 and 58 or between positions 60 and 61 in the amino acid sequence as shown in Seq ID No. 6.

Example 6Production of the plant transformation vector pTaSS1-as

In order to express a partial antisense-RNA to the isolated cDNA from wheat, a plant transformation vector was constructed on the basis of pUC19 as base plasmid. The plant transformation vector partially contains the cDNA insertion of the plasmid pTaSS1 in antisense-orientation. The expression is regulated by the ubiquitin-promoter. This promoter consists of the first untranslated exon and the first intron of the ubiquitin1 gene from maize (Christensen A. H. et al., Plant Molecular Biology 18 (1992), 675-689).

Parts of the polylinker and the NOS terminator are derived from the pAct.cas plasmid (CAMBIA, TG 0063; Cambia, GPO Box 3200, Canberra ACT 2601, Australia). Vector constructs with this terminator and constructs based on pAct1.cas are described in McElroy et al., (Molecular Breeding 1 (1995), 27-37).

In order to transform wheat the pTaSS1-as vector is used as described above.

Example 7Complementation of an E.coli mutant with a cDNA clone encoding a wheat soluble starch synthase

Enzymatic activity of the soluble starch synthase encoded by the cDNA clone TaSSS (Example 2) was analysed by complementation experiments using the E. coli mutant Hfr G6MD2 (M. Schwartz strain; CGSC # 5080; E.coli Genetic Stock Center, New Haven, USA) as host for gene expression. The E. coli mutant

shows a deletion of the *glg*-operon, encoding the bacterial ADP-glucose pyrophosphorylase (*glg* C), glycogen synthase (*glg* A) and branching enzyme (*glg* B). This mutation results in inability of glycogen synthesis through the ADP-glucose pathway. In addition, a deletion of the *mal* A operon prevents synthesis of linear α -1,4-glucans by the enzyme amylomaltase (*mal* Q).

The functionality of the soluble starch synthase was tested by cotransformation of the plasmids pTaSSSA188 and pACAG in the mutant G6MD2. The plasmid pTaSSSA188 comprise nucleotides 188-2239 of the 2239 bp cDNA sequence, which code for the soluble starch synthase. The cDNA is inserted as *Eco* RI/ *Xho* I fragment in the polylinker region of the pBluescript vector (Stratagene). This allows the N-terminus of the α -peptide of the beta-galactosidase encoded by the vector to be fused in frame with a part of the soluble starch synthase.

A successful complementation of the glycogen synthase (*glg* A) mutation in G6MD2 is dependent on expression of an ADP-glucose pyrophosphorylase activity, responsible for supply of ADP-glucose, the substrate for synthesis of α -1,4-glucans. Therefore, the plasmid pACAG (Abel G.J.W., (1995), Untersuchungen zur Funktion von Stärke-Synthasen in der Kartoffel (*Solanum tuberosum* L.), Dissertation, Freie Universität Berlin) comprising the coding region of the *glg* C locus isolated from the *E. coli* strain LCB 618 (Baecker et al., J. Biol. Chem, 258 (1983) 5084-5088) under control of the *lacZ* promoter was cotransformed. The encoded ADP-glucose pyrophosphorylase activity is less influenced by the activator fructose-1,6-bisphosphate and the inhibitor AMP resulting in sufficient supply of ADP-glucose.

Cells cotransformed with the constructs pTaSSSA188 and pACAG were plated out on LB-agar plates supplemented with 1% glucose,

1mM IPTG and 50 μ M diaminopimelate. The resulting colonies were stained by iodine steam. The transformed G6MD2 cells showed a blue-light brownish color in contrast to the yellowish colour of untransformed colonies, which indicates the ADP glucose: α -1,4-D-glucan 4- α -glucosyltransferase activity of the expressed fusion protein.

The system was checked by iodine staining of G6MD2 cells cotransformed with the constructs pACAG and pEc5.3. The plasmid pEc5.3 comprises a glycogen synthase (*glg A*) gene isolated from the E.coli strain DH5 α by PCR technologies (Abel G.J.W., (1995), Untersuchungen zur Funktion von Stärke-Synthasen in der Kartoffel (*Solanum tuberosum* L.), Dissertation, Freie Universität Berlin). The transformed cells showed a dark blue colour after staining with iodine, which indicates synthesis of α -1,4-glucans.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Hoechst Schering AgrEvo GmbH
- (B) STREET: Miraustrasse 54
- (C) CITY: Berlin
- (E) COUNTRY: DE
- (F) POSTAL CODE (ZIP): 13509

(ii) TITLE OF INVENTION: Nucleic acid molecules encoding enzymes from wheat which are involved in starch synthesis

(iii) NUMBER OF SEQUENCES: 9

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2239 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Triticum aestivum L.
- (B) STRAIN: cv. Florida
- (E) HAPLOTYPE: ca. 21 d Caryopses

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: cDNA library in pBluescript sk (-)
- (B) CLONE: TaSSS

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 3..2017

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CG ACG CAG CCG CCC CTG CCG GAC GCC GGC GTG GGG GAA CTC GCG CCC
 Thr Gln Pro Pro Leu Pro Asp Ala Gly Val Gly Glu Leu Ala Pro
 1 5 10 15

48

GAC CTC CTG CTC GAA GGG ATT GCT GAG GAT TCC ATC GAC AGC ATA ATT Asp Leu Leu Leu Glu Gly Ile Ala Glu Asp Ser Ile Asp Ser Ile Ile	95
20 25 30	
GTG GCT GCA AGT GAG CAG GAT TCT GAG ATC ATG GAT GCG AAT GAG CAA Val Ala Ala Ser Glu Gln Asp Ser Glu Ile Met Asp Ala Asn Glu Gln	143
35 40 45	
CCT CAA GCT AAA GTT ACA CGT AGC ATC GTG TTT GTG ACT GGT GAA GCT Pro Gln Ala Lys Val Thr Arg Ser Ile Val Phe Val Thr Gly Glu Ala	191
50 55 60	
GCT CCT TAT GCA AAG TCA GGG GGG TTG GGA GAT GTT TGT GGT TCG TTA Ala Pro Tyr Ala Lys Ser Gly Gly Leu Gly Asp Val Cys Gly Ser Leu	239
65 70 75	
CCA ATT GCT CTT GCT GCT CGT GGT CAC CGA GTG ATG GTT GTA ATG CCA Pro Ile Ala Leu Ala Ala Arg Gly His Arg Val Met Val Val Met Pro	287
80 85 90 95	
AGA TAC TTA AAT GGG TCC TCT GAT AAA AAC TAT GCA AAG GCA TTA TAC Arg Tyr Leu Asn Gly Ser Ser Asp Lys Asn Tyr Ala Lys Ala Leu Tyr	335
100 105 110	
ACT GCG AAG CAC ATT AAG ATT CCA TGC TTT GGG GGA TCA CAT GAA GTG Thr Ala Lys His Ile Lys Ile Pro Cys Phe Gly Gly Ser His Glu Val	383
115 120 125	
ACC TTT TTT CAT GAG TAT AGA GAC AAC GTC GAT TGG GTG TTT GTC GAT Thr Phe Phe His Glu Tyr Arg Asp Asn Val Asp Trp Val Phe Val Asp	431
130 135 140	
CAT CCG TCA TAT CAC AGA CCA GGA AGT TTA TAT GGA GAT AAT TTT GGT His Pro Ser Tyr His Arg Pro Gly Ser Leu Tyr Gly Asp Asn Phe Gly	479
145 150 155	
GCT TTT GGT GAT AAT CAG TTC AGA TAC ACA CTC CTT TGC TAT GCT GCA Ala Phe Gly Asp Asn Gln Phe Arg Tyr Thr Leu Leu Cys Tyr Ala Ala	527
160 165 170 175	
TGC GAG GCC CCA CTA ATC CTT GAA TTG GGA GGA TAT ATT TAT GGA CAG Cys Glu Ala Pro Leu Ile Leu Glu Leu Gly Gly Tyr Ile Tyr Gly Gln	575
180 185 190	
AAT TGC ATG TTT GTT GTG AAC GAT TGG CAT GCC AGC CTT GTG CCA GTC Asn Cys Met Phe Val Val Asn Asp Trp His Ala Ser Leu Val Pro Val	623
195 200 205	
CTT CTT GCT GCA AAA TAT AGA CCA TAC GGT GTT TAC AGA GAT TCC CGC Leu Leu Ala Ala Lys Tyr Arg Pro Tyr Gly Val Tyr Arg Asp Ser Arg	671
210 215 220	
AGC ACC CTT GTT ATA CAT AAT TTA GCA CAT CAG GGT GTG GAG CCT GCA Ser Thr Leu Val Ile His Asn Leu Ala His Gln Gly Val Glu Pro Ala	719
225 230 235	

AGT ACA TAT CCT GAT CTG GGA TTG CCT CCT GAA TGG TAT GGA GCT TTA	767
Ser Thr Tyr Pro Asp Leu Gly Leu Pro Pro Glu Trp Tyr Gly Ala Leu	
240 245 250 255	
GAA TGG GTA TTT CCA GAA TGG GCA AGG AGG CAT GCC CTT GAC AAG GGT	815
Glu Trp Val Phe Pro Glu Trp Ala Arg Arg His Ala Leu Asp Lys Gly	
260 265 270	
GAG GCA GTT AAC TTT TTG AAA GGA GCA GTT GTG ACA GCA GAT CGG ATT	863
Glu Ala Val Asn Phe Leu Lys Gly Ala Val Val Thr Ala Asp Arg Ile	
275 280 285	
GTG ACC GTC AGT CAG GGT TAT TCA TGG GAG GTC ACA ACT GCT GAA GGT	911
Val Thr Val Ser Gln Gly Tyr Ser Trp Glu Val Thr Thr Ala Glu Gly	
290 295 300	
GGA CAG GGC CTC AAT GAG CTC TTA AGC TCC CGA AAA AGT GTA TTG AAT	959
Gly Gln Gly Leu Asn Glu Leu Leu Ser Ser Arg Lys Ser Val Leu Asn	
305 310 315	
GGA ATT GTA AAT GGA ATT GAC ATT AAT GAT TGG AAC CCC ACC ACA GAC	1007
Gly Ile Val Asn Gly Ile Asp Ile Asn Asp Trp Asn Pro Thr Thr Asp	
320 325 330 335	
AAG TGT CTC CCT CAT CAT TAT TCT GTC GAT GAC CTC TCT GGA AAG GCC	1055
Lys Cys Leu Pro His His Tyr Ser Val Asp Asp Leu Ser Gly Lys Ala	
340 345 350	
AAA TGT AAA GCT GAA TTG CAG AAG GAG TTG GGT TTA CCT GTA AGG GAG	1103
Lys Cys Lys Ala Glu Leu Gln Lys Glu Leu Gly Leu Pro Val Arg Glu	
355 360 365	
GAT GTT CCT CTG ATT GGC TTT ATT GGA AGA CTG GAT TAC CAG AAA GGC	1151
Asp Val Pro Leu Ile Gly Phe Ile Gly Arg Leu Asp Tyr Gln Lys Gly	
370 375 380	
ATT GAT CTC ATT AAA ATG GCC ATT CCA GAG CTC ATG AGG GAG GAC GTG	1199
Ile Asp Leu Ile Lys Met Ala Ile Pro Glu Leu Met Arg Glu Asp Val	
385 390 395	
CAA TTT GTC ATG CTT GGA TCT GGG GAT CCA ATT TTT GAA GGC TGG ATG	1247
Gln Phe Val Met Leu Gly Ser Gly Asp Pro Ile Phe Glu Gly Trp Met	
400 405 410 415	
AGA TCT ACC GAG TCG AGT TAC AAG GAT AAA TTC CGT GGA TGG GTT GGA	1295
Arg Ser Thr Glu Ser Ser Tyr Lys Asp Lys Phe Arg Gly Trp Val Gly	
420 425 430	
TTT AGT GTT CCA GTT TCC CAC AGA ATA ACT GCA GGT TGC GAT ATA TTG	1343
Phe Ser Val Pro Val Ser His Arg Ile Thr Ala Gly Cys Asp Ile Leu	
435 440 445	
TTA ATG CCA TCG AGA TTT GAA CCT TGC GGT CTT AAT CAG CTA TAT GCT	1391
Leu Met Pro Ser Arg Phe Glu Pro Cys Gly Leu Asn Gln Leu Tyr Ala	
450 455 460	

ATG CAA TAT GGT ACA GTT CCT GTA GTT CAT GGA ACT GGG GGC CTC CGA Met Gln Tyr Gly Thr Val Pro Val Val His Gly Thr Gly Gly Leu Arg 465 470 475	1439
GAC ACA GTC GAG ACC TTC AAC CCT TTT GGT GCA AAA GGA GAG GAG GGT Asp Thr Val Glu Thr Phe Asn Pro Phe Gly Ala Lys Gly Glu Glu Gly 480 485 490 495	1487
ACA GGG TGG GCG TTC TCA CCG CTA ACC GTG GAC AAG ATG TTG TGG GCA Thr Gly Trp Ala Phe Ser Pro Leu Thr Val Asp Lys Met Leu Trp Ala 500 505 510	1535
TTG CGA ACC GCG ATG TCG ACA TTC AGG GAG CAC AAG CCG TCC TGG GAG Leu Arg Thr Ala Met Ser Thr Phe Arg Glu His Lys Pro Ser Trp Glu 515 520 525	1583
GGG CTC ATG AAG CGA GGC ATG ACG AAA GAC CAT ACG TGG GAC CAT GCC Gly Leu Met Lys Arg Gly Met Thr Lys Asp His Thr Trp Asp His Ala 530 535 540	1631
CCG AGC AGT ACG AGC AGA TCT TCG AGT GGG CCT TCG TGG ACC AAC CCT Pro Ser Ser Thr Ser Arg Ser Ser Ser Gly Pro Ser Trp Thr Asn Pro 545 550 555	1679
ACG TCA TGT AGA CGG GGA CTG GGG AGG TCC AAG TGC GAG TCT CCT TCA Thr Ser Cys Arg Arg Gly Leu Gly Arg Ser Lys Cys Glu Ser Pro Ser 560 565 570 575	1727
GCT CTG AAG ACA TCC TCT TCA TCC TTC CGC GGC CCG GAA GGA TAC CCC Ala Leu Lys Thr Ser Ser Ser Ser Phe Arg Gly Pro Glu Gly Tyr Pro 580 585 590	1775
TGT ACA TTG CGT TGT CCT GCT ACA GTA GAG TCG CAA TGC GCC TGC TTG Cys Thr Leu Arg Cys Pro Ala Thr Val Glu Ser Gln Cys Ala Cys Leu 595 600 605	1823
CTT TGG TTC GCC GGT TCG AGA ACA TAT GAC GGC TGT GCT GCT GCG GCG Leu Trp Phe Ala Gly Ser Arg Thr Tyr Asp Gly Cys Ala Ala Ala Ala 610 615 620	1871
GTG ACA GCT TCG GGT GGA CGA CAG TTA CAG TTT TGG GGA ATA AGG AAG Val Thr Ala Ser Gly Gly Arg Gln Leu Gln Phe Trp Gly Ile Arg Lys 625 630 635	1919
GGA TGT GCT GCA GGA TGG TTA ACA GCA AAG CAC CAC TCA GAT GGC AGC Gly Cys Ala Ala Gly Trp Leu Thr Ala Lys His His Ser Asp Gly Ser 640 645 650 655	1967
CTC TCT GTC CGT GTT ACA GCT GAA ATC AGA AAC CAA CTG GTG ACT CTT TA Leu Ser Val Arg Val Thr Ala Glu Ile Arg Asn Gln Leu Val Thr Leu 660 665 670	2017
GCCTTAGTGA TTGTGAAGTT TGTTCCTTC TGTGTATGTT GTCTTGTCTT TAGCTGACAA	2077
ATATTTGACC TGTTGGAGAA TTTTATCTTT GCTGCTGTTT TTTTITAATC AAAAGAGGGG	2137
GTTTCCTCCG ATTCATTAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA	2197

AAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AA

2239

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 671 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Thr Gln Pro Pro Leu Pro Asp Ala Gly Val Gly Glu Leu Ala Pro Asp
 1 5 10 15
 Leu Leu Leu Glu Gly Ile Ala Glu Asp Ser Ile Asp Ser Ile Ile Val
 20 25 30
 Ala Ala Ser Glu Gln Asp Ser Glu Ile Met Asp Ala Asn Glu Gln Pro
 35 40 45
 Gln Ala Lys Val Thr Arg Ser Ile Val Phe Val Thr Gly Glu Ala Ala
 50 55 60
 Pro Tyr Ala Lys Ser Gly Gly Leu Gly Asp Val Cys Gly Ser Leu Pro
 65 70 75 80
 Ile Ala Leu Ala Ala Arg Gly His Arg Val Met Val Val Met Pro Arg
 85 90 95
 Tyr Leu Asn Gly Ser Ser Asp Lys Asn Tyr Ala Lys Ala Leu Tyr Thr
 100 105 110
 Ala Lys His Ile Lys Ile Pro Cys Phe Gly Gly Ser His Glu Val Thr
 115 120 125
 Phe Phe His Glu Tyr Arg Asp Asn Val Asp Trp Val Phe Val Asp His
 130 135 140
 Pro Ser Tyr His Arg Pro Gly Ser Leu Tyr Gly Asp Asn Phe Gly Ala
 145 150 155 160
 Phe Gly Asp Asn Gln Phe Arg Tyr Thr Leu Leu Cys Tyr Ala Ala Cys
 165 170 175
 Glu Ala Pro Leu Ile Leu Glu Leu Gly Gly Tyr Ile Tyr Gly Gln Asn
 180 185 190
 Cys Met Phe Val Val Asn Asp Trp His Ala Ser Leu Val Pro Val Leu
 195 200 205
 Leu Ala Ala Lys Tyr Arg Pro Tyr Gly Val Tyr Arg Asp Ser Arg Ser
 210 215 220
 Thr Leu Val Ile His Asn Leu Ala His Gln Gly Val Glu Pro Ala Ser
 225 230 235 240

Thr Tyr Pro Asp Leu Gly Leu Pro Pro Glu Trp Tyr Gly Ala Leu Glu
 245 250 255
 Trp Val Phe Pro Glu Trp Ala Arg Arg His Ala Leu Asp Lys Gly Glu
 260 265 270
 Ala Val Asn Phe Leu Lys Gly Ala Val Val Thr Ala Asp Arg Ile Val
 275 280 285
 Thr Val Ser Gln Gly Tyr Ser Trp Glu Val Thr Thr Ala Glu Gly Gly
 290 295 300
 Gln Gly Leu Asn Glu Leu Leu Ser Ser Arg Lys Ser Val Leu Asn Gly
 305 310 315 320
 Ile Val Asn Gly Ile Asp Ile Asn Asp Trp Asn Pro Thr Thr Asp Lys
 325 330 335
 Cys Leu Pro His His Tyr Ser Val Asp Asp Leu Ser Gly Lys Ala Lys
 340 345 350
 Cys Lys Ala Glu Leu Gln Lys Glu Leu Gly Leu Pro Val Arg Glu Asp
 355 360 365
 Val Pro Leu Ile Gly Phe Ile Gly Arg Leu Asp Tyr Gln Lys Gly Ile
 370 375 380
 Asp Leu Ile Lys Met Ala Ile Pro Glu Leu Met Arg Glu Asp Val Gln
 385 390 395 400
 Phe Val Met Leu Gly Ser Gly Asp Pro Ile Phe Glu Gly Trp Met Arg
 405 410 415
 Ser Thr Glu Ser Ser Tyr Lys Asp Lys Phe Arg Gly Trp Val Gly Phe
 420 425 430
 Ser Val Pro Val Ser His Arg Ile Thr Ala Gly Cys Asp Ile Leu Leu
 435 440 445
 Met Pro Ser Arg Phe Glu Pro Cys Gly Leu Asn Gln Leu Tyr Ala Met
 450 455 460
 Gln Tyr Gly Thr Val Pro Val Val His Gly Thr Gly Gly Leu Arg Asp
 465 470 475 480
 Thr Val Glu Thr Phe Asn Pro Phe Gly Ala Lys Gly Glu Glu Gly Thr
 485 490 495
 Gly Trp Ala Phe Ser Pro Leu Thr Val Asp Lys Met Leu Trp Ala Leu
 500 505 510
 Arg Thr Ala Met Ser Thr Phe Arg Glu His Lys Pro Ser Trp Glu Gly
 515 520 525
 Leu Met Lys Arg Gly Met Thr Lys Asp His Thr Trp Asp His Ala Pro
 530 535 540

53

Ser Ser Thr Ser Arg Ser Ser Ser Gly Pro Ser Trp Thr A ro Thr
 545 550 555 560

Ser Cys Arg Arg Gly Leu Gly Arg Ser Lys Cys Glu Ser Pro Ser Ala
 565 570 575

Leu Lys Thr Ser Ser Ser Ser Phe Arg Gly Pro Glu Gly Tyr Pro Cys
 580 585 590

Thr Leu Arg Cys Pro Ala Thr Val Glu Ser Gln Cys Ala Cys Leu Leu
 595 600 605

Trp Phe Ala Gly Ser Arg Thr Tyr Asp Gly Cys Ala Ala Ala Ala Val
 610 615 620

Thr Ala Ser Gly Gly Arg Gln Leu Gln Phe Trp Gly Ile Arg Lys Gly
 625 630 635 640

Cys Ala Ala Gly Trp Leu Thr Ala Lys His His Ser Asp Gly Ser Leu
 645 650 655

Ser Val Arg Val Thr Ala Glu Ile Arg Asn Gln Leu Val Thr Leu
 660 665 670

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "Oligonucleotide"

(iii) HYPOTHETICAL: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

ACAGGATCCT GTGCTATGCG GCGTGTGAAG

30

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "oligonucleotide"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

TTGGGATCCG CAATGCCCAC AGCATTTC TC

32

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2825 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Triticum aestivum L.
- (B) STRAIN: cv. Florida
- (F) TISSUE TYPE: ca. 21 d Caryopses

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: cDNA library in pBluescript sk (-)
- (B) CLONE: pTASS1

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION:162..2559

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CTTCGGCCTG ACCCCGTTTCG TTTACCCCCA CACAGAGCAC ACTCCAGTCC AGTCCAGCCC	60
ACTGCCACCG CGCTACTCTC CACTCCCACT GCCACCACCT CCGCCTGCGC CGCGCTCTGG	120
GCGGACCAAC CCGCGAACCG TACCATCTCC CGCCCCGATC C ATG TCG TCG GCG	173
Met Ser Ser Ala	675
GTC GCG TCC GCC GCA TCC TTC CTC GCG CTC GCG TCA GCC TCC CCC GGG	221
Val Ala Ser Ala Ala Ser Phe Leu Ala Leu Ala Ser Ala Ser Pro Gly	
680 685 690	
AGA TCA CGC AGG CGG GCG AGG GTG AGC GCG CAG CCA CCC CAC GCC GGG	269
Arg Ser Arg Arg Arg Ala Arg Val Ser Ala Gln Pro Pro His Ala Gly	
695 700 705	
GCC GGC AGG TTG CAC TGG CCG CCG TGG CCG CCG CAG CGC ACG GCT CGC	317
Ala Gly Arg Leu His Trp Pro Pro Trp Pro Pro Gln Arg Thr Ala Arg	
710 715 720	

GAC	GGA	GCT	GTG	GCG	GCG	CTC	GCC	GCC	GGG	AAG	AAG	GAC	GCG	GGG	ATC	365
Asp	Gly	Ala	Val	Ala	Ala	Leu	Ala	Ala	Gly	Lys	Lys	Asp	Ala	Gly	Ile	
	725					730					735					
GAC	GAC	GCC	GCC	GCG	TCC	GTG	AGG	CAG	CCC	CGC	GCA	CTC	CGC	GGT	GGC	413
Asp	Asp	Ala	Ala	Ala	Ser	Val	Arg	Gln	Pro	Arg	Ala	Leu	Arg	Gly	Gly	
	740				745				750					755		
GCC	GCC	ACC	AAG	GTC	GCG	GAG	CGA	AGG	GAT	CCC	GTC	AAG	ACG	CTC	GAC	461
Ala	Ala	Thr	Lys	Val	Ala	Glu	Arg	Arg	Asp	Pro	Val	Lys	Thr	Leu	Asp	
				760					765					770		
CGC	GAC	GCC	GCG	GAA	GGC	GGC	GGG	CCG	TCC	CCG	CCG	GCA	GCG	AGG	CAG	509
Arg	Asp	Ala	Ala	Glu	Gly	Gly	Gly	Pro	Ser	Pro	Pro	Ala	Ala	Arg	Gln	
			775					780						785		
GAC	GCC	GCC	CGT	CCG	CCG	AGT	ATG	AAC	GGC	ATG	CCG	GTG	AAC	GGC	GAG	557
Asp	Ala	Ala	Arg	Pro	Pro	Ser	Met	Asn	Gly	Met	Pro	Val	Asn	Gly	Glu	
		790					795					800				
AAC	AAA	TCT	ACC	GGC	GGC	GGC	GGC	GCG	ACT	AAA	GAC	AGC	GGG	CTG	CCC	605
Asn	Lys	Ser	Thr	Gly	Gly	Gly	Gly	Ala	Thr	Lys	Asp	Ser	Gly	Leu	Pro	
	805						810				815					
ACG	CCC	GCA	CGC	GCG	CCC	CAT	CCG	TCG	ACC	CAG	AAC	AGA	GCA	CCG	GTG	653
Thr	Pro	Ala	Arg	Ala	Pro	His	Pro	Ser	Thr	Gln	Asn	Arg	Ala	Pro	Val	
	820				825					830					835	
AAC	GGT	GAA	AAC	AAA	GCT	AAC	GTC	GCC	TCG	CCG	CCG	ACG	AGC	ATA	GCC	701
Asn	Gly	Glu	Asn	Lys	Ala	Asn	Val	Ala	Ser	Pro	Pro	Thr	Ser	Ile	Ala	
				840					845					850		
GAG	GCC	GCG	GCT	TCG	GAT	TCC	GCA	GCT	ACC	ATT	TCC	ATC	AGC	GAC	AAG	749
Glu	Ala	Ala	Ala	Ser	Asp	Ser	Ala	Ala	Thr	Ile	Ser	Ile	Ser	Asp	Lys	
			855						860					865		
GCG	CCG	GAG	TCC	GTT	GTC	CCA	GCT	GAG	AAG	ACG	CCG	CCG	TCG	TCC	GGC	797
Ala	Pro	Glu	Ser	Val	Val	Pro	Ala	Glu	Lys	Thr	Pro	Pro	Ser	Ser	Gly	
		870						875					880			
TCA	AAT	TTC	GAG	TCC	TCG	GCC	TCT	GCT	CCC	GGG	TCT	GAC	ACT	GTC	AGC	845
Ser	Asn	Phe	Glu	Ser	Ser	Ala	Ser	Ala	Pro	Gly	Ser	Asp	Thr	Val	Ser	
		885				890						895				
GAC	GTG	GAA	CAA	GAA	CTG	AAG	AAG	GGT	GCG	GTC	GTT	GTC	GAA	GAA	GCT	893
Asp	Val	Glu	Gln	Glu	Leu	Lys	Lys	Gly	Ala	Val	Val	Val	Glu	Glu	Ala	
	900				905				910					915		
CCA	AAG	CCA	AAG	GCT	CTT	TCG	CCG	CCT	GCA	GCC	CCC	GCT	GTA	CAA	GAA	941
Pro	Lys	Pro	Lys	Ala	Leu	Ser	Pro	Pro	Ala	Ala	Pro	Ala	Val	Gln	Glu	
				920					925					930		
GAC	CTT	TGG	GAT	TTC	AAG	AAA	TAC	ATT	GGT	TTC	GAG	GAG	CCC	GTG	GAG	989
Asp	Leu	Trp	Asp	Phe	Lys	Lys	Tyr	Ile	Gly	Phe	Glu	Glu	Pro	Val	Glu	
			935					940						945		

GCC AAG GAT GAT GGC CGG GCT GTC GCA GAT GAT GCG GGC TCC TTT GAA Ala Lys Asp Asp Gly Arg Ala Val Ala Asp Asp Ala Gly Ser Phe Glu 950 955 960	1037
CAC CAC CAG AAT CAC GAC TCC GGA CCT TTG GCA GGG GAG AAT GTC ATG His His Gln Asn His Asp Ser Gly Pro Leu Ala Gly Glu Asn Val Met 965 970 975	1085
AAC GTG GTC GTC GTG GCT GCT GAG TGT TCT CCC TGG TGC AAA ACA GGT Asn Val Val Val Val Ala Ala Glu Cys Ser Pro Trp Cys Lys Thr Gly 980 985 990 995	1133
GGT CTG GGA GAT GTT GCG GGT GCT CTG CCC AAG GCT TTG GCA AAG AGA Gly Leu Gly Asp Val Ala Gly Ala Leu Pro Lys Ala Leu Ala Lys Arg 1000 1005 1010	1181
GGA CAT CGT GTT ATG GTT GTG GTA CCA AGG TAT GGG GAC TAT GAA GAA Gly His Arg Val Met Val Val Val Pro Arg Tyr Gly Asp Tyr Glu Glu 1015 1020 1025	1229
GCC TAC GAT GTC GGA GTC CGA AAA TAC TAC AAG GCT GCT GGA CAG GAT Ala Tyr Asp Val Gly Val Arg Lys Tyr Tyr Lys Ala Ala Gly Gln Asp 1030 1035 1040	1277
ATG GAA GTG AAT TAT TTC CAT GCT TAT ATC GAT GGA GTT GAT TTT GTG Met Glu Val Asn Tyr Phe His Ala Tyr Ile Asp Gly Val Asp Phe Val 1045 1050 1055	1325
TTC ATT GAC GCT CCT CTC TTC CGA CAC CGT CAG GAA GAC ATT TAT GGG Phe Ile Asp Ala Pro Leu Phe Arg His Arg Gln Glu Asp Ile Tyr Gly 1060 1065 1070 1075	1373
GGC AGC AGA CAG GAA ATT ATG AAG CGC ATG ATT TTG TTC TGC AAG GCC Gly Ser Arg Gln Glu Ile Met Lys Arg Met Ile Leu Phe Cys Lys Ala 1080 1085 1090	1421
GCT GTT GAG GTT CCA TGG CAC GTT CCA TGC GGC GGT GTC CCT TAT GGG Ala Val Glu Val Pro Trp His Val Pro Cys Gly Gly Val Pro Tyr Gly 1095 1100 1105	1469
GAT GGA AAT CTG GTG TTT ATT GCA AAT GAT TGG CAC ACG GCA CTC CTG Asp Gly Asn Leu Val Phe Ile Ala Asn Asp Trp His Thr Ala Leu Leu 1110 1115 1120	1517
CCT GTC TAT CTG AAA GCA TAT TAC AGG GAC CAT GGT TTG ATG CAG TAC Pro Val Tyr Leu Lys Ala Tyr Tyr Arg Asp His Gly Leu Met Gln Tyr 1125 1130 1135	1565
ACT CGG TCC ATT ATG GTG ATA CAT AAC ATC GCT CAC CAG GGC CGT GGC Thr Arg Ser Ile Met Val Ile His Asn Ile Ala His Gln Gly Arg Gly 1140 1145 1150 1155	1613
CCT GTA GAT GAA TTC CCG TTC ACC GAG TTG CCT GAG CAC TAC CTG GAA Pro Val Asp Glu Phe Pro Phe Thr Glu Leu Pro Glu His Tyr Leu Glu 1160 1165 1170	1661

CAC TTC AGA CTG TAC GAC CCC GTG GGT GGT GAA CAC GCC AAC TAC TTC	1709
His Phe Arg Leu Tyr Asp Pro Val Gly Gly Glu His Ala Asn Tyr Phe	
1175 1180 1185	
GCC GCC GGC CTG AAG ATG GCG GAC CAG GTT GTC GTG GTG AGC CCC GGG	1757
Ala Ala Gly Leu Lys Met Ala Asp Gln Val Val Val Val Ser Pro Gly	
1190 1195 1200	
TAC CTG TGG GAG CTG AAG ACG GTG GAG GGC GGC TGG GGG CTT CAC GAC	1805
Tyr Leu Trp Glu Leu Lys Thr Val Glu Gly Gly Trp Gly Leu His Asp	
1205 1210 1215	
ATC ATA CGG CAG AAC GAC TGG AAG ACC CGC GGC ATC GTC AAC GGC ATC	1853
Ile Ile Arg Gln Asn Asp Trp Lys Thr Arg Gly Ile Val Asn Gly Ile	
1220 1225 1230 1235	
GAC AAC ATG GAG TGG AAC CCC GAG GTG GAC GCC CAC CTC AAG TCG GAC	1901
Asp Asn Met Glu Trp Asn Pro Glu Val Asp Ala His Leu Lys Ser Asp	
1240 1245 1250	
GGC TAC ACC AAC TTC TCC CTG AGG ACG CTG GAC TCC GGC AAG CGG CAG	1949
Gly Tyr Thr Asn Phe Ser Leu Arg Thr Leu Asp Ser Gly Lys Arg Gln	
1255 1260 1265	
TGC AAG GAG GCC CTG CAG CGC GAG CTG GGC CTG CAG GTC CGC GCC GAC	1997
Cys Lys Glu Ala Leu Gln Arg Glu Leu Gly Leu Gln Val Arg Ala Asp	
1270 1275 1280	
GTG CCG CTG CTC GGC TTC ATC GGC CGC CTG GAC GGG CAG AAG GGC GTG	2045
Val Pro Leu Leu Gly Phe Ile Gly Arg Leu Asp Gly Gln Lys Gly Val	
1285 1290 1295	
GAG ATC ATC GCG GAC GCC ATG CCC TGG ATC GTG AGC CAG GAC GTG CAG	2093
Glu Ile Ile Ala Asp Ala Met Pro Trp Ile Val Ser Gln Asp Val Gln	
1300 1305 1310 1315	
CTG GTG ATG CTG GGC ACC GGG CGC CAC GAC CTG GAG AGC ATG CTG CAG	2141
Leu Val Met Leu Gly Thr Gly Arg His Asp Leu Glu Ser Met Leu Gln	
1320 1325 1330	
CAC TTC GAG CGG GAG CAC CAC GAC AAG GTG CGC GGG TGG GTG GGG TTC	2189
His Phe Glu Arg Glu His His Asp Lys Val Arg Gly Trp Val Gly Phe	
1335 1340 1345	
TCC GTG CGC CTG GCG CAC CGG ATC ACG GCG GGG GCG GAC GCG CTC CTC	2237
Ser Val Arg Leu Ala His Arg Ile Thr Ala Gly Ala Asp Ala Leu Leu	
1350 1355 1360	
ATG CCC TCC CGG TTC GAG CCG TGC GGG CTG AAC CAG CTC TAC GCC ATG	2285
Met Pro Ser Arg Phe Glu Pro Cys Gly Leu Asn Gln Leu Tyr Ala Met	
1365 1370 1375	
GCC TAC GGC ACC GTC CCC GTC GTG CAC GCC GTC GGC GGC CTC AGG GAC	2333
Ala Tyr Gly Thr Val Pro Val Val His Ala Val Gly Gly Leu Arg Asp	
1380 1385 1390 1395	

ACC GTG CCG CCG TTC GAC CCC TTC AAC CAC TCC GGG CTC GGG TGG ACG	2381
Thr Val Pro Pro Phe Asp Pro Phe Asn His Ser Gly Leu Gly Trp Thr	
1400 1405 1410	
TTC GAC CGC GCC GAG GCG CAC AAG CTG ATC GAG GCG CTC GGG CAC TGC	2429
Phe Asp Arg Ala Glu Ala His Lys Leu Ile Glu Ala Leu Gly His Cys	
1415 1420 1425	
CTC CGC ACC TAC CGA GAC TTC AAG GAG AGC TGG AGG GCC CTC CAG GAG	2477
Leu Arg Thr Tyr Arg Asp Phe Lys Glu Ser Trp Arg Ala Leu Gln Glu	
1430 1435 1440	
CGC GGC ATG TCG CAG GAC TTC AGC TGG GAG CAC GCC GCC AAG CTC TAC	2525
Arg Gly Met Ser Gln Asp Phe Ser Trp Glu His Ala Ala Lys Leu Tyr	
1445 1450 1455	
GAG GAC GTC CTC GTC AAG GCC AAG TAC CAG TGG T GAACGCTAGC	2569
Glu Asp Val Leu Val Lys Ala Lys Tyr Gln Trp	
1460 1465 1470	
TGCTAGCCGC TCCAGCCCCG CATGCGTGCA TGACAGGATG GAACTGCATT GCGCACGCAG	2629
GAAAGTGCCA TGGAGCGCCG GCATCCGCGA AGTACAGTGA CATGAGGTGT GTGTGGTTGA	2689
GACGCTGATT CCAATCCGGC CCGTAGCAGA GTAGAGCGGA GGTATATGGG AATCTTAACT	2749
TGGTATTGTA ATTTGTTATG TTGTGTGCAT TATTACAATG TTGTTACTTA TTCTTGTTAA	2809
AAAAAAAAA AAAAAA	2825

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 799 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Met Ser Ser Ala Val Ala Ser Ala Ala Ser Phe Leu Ala Leu Ala Ser	
1 5 10 15	
Ala Ser Pro Gly Arg Ser Arg Arg Arg Ala Arg Val Ser Ala Gln Pro	
20 25 30	
Pro His Ala Gly Ala Gly Arg Leu His Trp Pro Pro Trp Pro Pro Gln	
35 40 45	
Arg Thr Ala Arg Asp Gly Ala Val Ala Ala Leu Ala Ala Gly Lys Lys	
50 55 60	
Asp Ala Gly Ile Asp Asp Ala Ala Ala Ser Val Arg Gln Pro Arg Ala	
65 70 75 80	

SUBSTITUTE SHEET (RULE 26)

60

Val Asp Phe Val Phe	Ile Asp Ala Pro Leu Phe Arg His Arg Gln Glu	
385	390	395 400
Asp Ile Tyr Gly Gly Ser Arg Gln Glu Ile Met Lys Arg Met Ile Leu		
	405	410 415
Phe Cys Lys Ala Ala Val Glu Val Pro Trp His Val Pro Cys Gly Gly		
	420	425 430
Val Pro Tyr Gly Asp Gly Asn Leu Val Phe Ile Ala Asn Asp Trp His		
	435	440 445
Thr Ala Leu Leu Pro Val Tyr Leu Lys Ala Tyr Tyr Arg Asp His Gly		
	450	455 460
Leu Met Gln Tyr Thr Arg Ser Ile Met Val Ile His Asn Ile Ala His		
	465	470 475 480
Gln Gly Arg Gly Pro Val Asp Glu Phe Pro Phe Thr Glu Leu Pro Glu		
	485	490 495
His Tyr Leu Glu His Phe Arg Leu Tyr Asp Pro Val Gly Gly Glu His		
	500	505 510
Ala Asn Tyr Phe Ala Ala Gly Leu Lys Met Ala Asp Gln Val Val Val		
	515	520 525
Val Ser Pro Gly Tyr Leu Trp Glu Leu Lys Thr Val Glu Gly Gly Trp		
	530	535 540
Gly Leu His Asp Ile Ile Arg Gln Asn Asp Trp Lys Thr Arg Gly Ile		
	545	550 555 560
Val Asn Gly Ile Asp Asn Met Glu Trp Asn Pro Glu Val Asp Ala His		
	565	570 575
Leu Lys Ser Asp Gly Tyr Thr Asn Phe Ser Leu Arg Thr Leu Asp Ser		
	580	585 590
Gly Lys Arg Gln Cys Lys Glu Ala Leu Gln Arg Glu Leu Gly Leu Gln		
	595	600 605
Val Arg Ala Asp Val Pro Leu Leu Gly Phe Ile Gly Arg Leu Asp Gly		
	610	615 620
Gln Lys Gly Val Glu Ile Ile Ala Asp Ala Met Pro Trp Ile Val Ser		
	625	630 635 640
Gln Asp Val Gln Leu Val Met Leu Gly Thr Gly Arg His Asp Leu Glu		
	645	650 655
Ser Met Leu Gln His Phe Glu Arg Glu His His Asp Lys Val Arg Gly		
	660	665 670
Trp Val Gly Phe Ser Val Arg Leu Ala His Arg Ile Thr Ala Gly Ala		
	675	680 685

Asp Ala Leu Leu Met Pro Ser Arg Phe Glu Pro Cys Gly Leu Asn Gln
 690 695 700
 Leu Tyr Ala Met Ala Tyr Gly Thr Val Pro Val Val His Ala Val Gly
 705 710 715 720
 Gly Leu Arg Asp Thr Val Pro Pro Phe Asp Pro Phe Asn His Ser Gly
 725 730 735
 Leu Gly Trp Thr Phe Asp Arg Ala Glu Ala His Lys Leu Ile Glu Ala
 740 745 750
 Leu Gly His Cys Leu Arg Thr Tyr Arg Asp Phe Lys Glu Ser Trp Arg
 755 760 765
 Ala Leu Gln Glu Arg Gly Met Ser Gln Asp Phe Ser Trp Glu His Ala
 770 775 780
 Ala Lys Leu Tyr Glu Asp Val Leu Val Lys Ala Lys Tyr Gln Trp
 785 790 795

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Pro Trp Ser Lys Thr Gly Gly Leu Gly Asp Val Cys
 1 5 10

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 13 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Pro	Ser	Arg	Phe	Glu	Pro	Cys	Gly	Leu	Asn	Gln	Leu	Tyr
1				5					10			

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Gly	Thr	Gly	Gly	Leu	Arg	Asp	Thr	Val	Glu	Asn	Cys
1				5					10		

Claims

1. A nucleic acid molecule encoding a starch synthase from wheat selected from the group consisting of
 - (a) nucleic acid molecules encoding a protein comprising the amino acid sequence indicated under Seq ID No.2;
 - (b) nucleic acid molecules comprising the nucleotide sequence indicated under Seq ID No. 1 or a corresponding ribonucleotide sequence;
 - (c) nucleic acid molecules which hybridize to the nucleic acid molecules mentioned under (a) or (b) and encode a soluble starch synthase;
 - (d) nucleic acid molecules, the nucleotide sequence of which deviates from the sequence of the nucleic acid molecules mentioned under (a), (b) or (c) due to the degeneracy of the genetic code;
 - (e) nucleic acid molecules encoding a protein which comprises the amino acid sequence indicated under Seq ID No. 6;
 - (f) nucleic acid molecules comprising the nucleotide sequence indicated under Seq ID No. 5 or a corresponding ribonucleotide sequence;
 - (g) nucleic acid molecules hybridizing to the nucleic acid molecules mentioned under (e) or (f); and
 - (h) nucleic acid molecules, the nucleotide sequence of which deviates from the sequence of the molecules mentioned under (e) to (g) due to the degeneracy of the genetic code.
2. The nucleic acid molecule of claim 1 which is a DNA molecule.

3. The DNA molecule of claim 2 which is a cDNA molecule.
4. The nucleic acid molecule of claim 1 which is an RNA molecule.
5. A nucleic acid molecule which specifically hybridizes to a strand of a nucleic acid molecule of any one of claims 1 to 4.
6. The nucleic acid molecule of claim 5 which is an oligonucleotide with a length of at least 15 nucleotides.
7. A vector containing a nucleic acid molecule of any one of claims 1 to 4.
8. The vector of claim 7, wherein the nucleic acid molecule is linked in sense-orientation to regulatory elements which ensure the transcription and the synthesis of a translatable RNA in procaryotic or eucaryotic cells.
9. A host cell which is transformed with a nucleic acid molecule of any one of claims 1 to 4 or with a vector of claim 7 or 8, or which is derived from such a cell.
10. A protein encoded by a nucleic acid molecule of any one of claims 1 to 4.
11. A method for the production of a protein according to claim 10 in which a host cell according to claim 9 is cultivated under conditions that allow for the synthesis of the protein and in which the protein is isolated from the cultivated cells and/or the culture medium.

12. A transgenic plant cell transformed with a nucleic acid molecule of any one of claims 1 to 4 or with a vector of claim 7 or 8, or derived from such a cell, wherein the nucleic acid molecule encoding a soluble starch synthase from wheat is subject to the control of regulatory elements allowing for the transcription of a translatable mRNA in plant cells.
13. A plant comprising plant cells of claim 12.
14. The plant of claim 13 which is a useful plant.
15. The plant of claim 14 which is a starch-storing plant.
16. The plant of claim 15 which is a wheat plant.
17. Propagation material of a plant of any one of claims 13 to 16, containing plant cells of claim 12.
18. Starch obtainable from a plant of any one of claims 13 to 16 or from propagation material of claim 17.
19. A transgenic plant cell which is characterized in that the activity of a protein of claim 10 is reduced in this plant cell.
20. The plant cell according to claim 19 wherein the reduction of the activity in this cell is achieved by the expression of an antisense-RNA to transcripts of a DNA molecule of claim 1.

21. A plant containing plant cells of claim 19 or 20.
22. The plant of claim 21 which is a useful plant.
23. The plant of claim 22 which is a starch-storing plant.
24. The plant of claim 23 which is a wheat plant.
25. Propagation material of a plant of any one of claims 21 to 24 containing cells of claim 19 or 20.
26. Starch obtainable from plants of any one of claims 21 to 24 or from propagation material of claim 25.
27. Use of the starch of claims 18 or 26 for the production of foodstuffs.
28. The use of claim 27, wherein the said foodstuffs are bakery-goods or pasta.
29. Use of the starch of claim 18 or 26 for the production of packaging material or disposable goods.

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/54 C12N15/82 C12N15/11 C12N9/10 C12N5/10
C08B39/00 A01H5/00 A23L1/0522

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C08B A01H A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	AINSWORTH C. ET AL.: "Expression, organisation and structure of the genes encoding the waxy protein (granule-bound starch synthase) in wheat" PLANT MOLECULAR BIOLOGY, vol. 22, no. 1, April 1993, pages 67-82, XP002043125 cited in the application see the whole document & "AC X57233" EMBL DATABASE, 2 August 1991, HEIDELBERG, see the whole document ---	1-11
X	DE 44 41 408 A (INST GENBIOLOGISCHE FORSCHUNG) 15 May 1996 see the whole document ---	18,26-29
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

10 October 1997

Date of mailing of the international search report

2 1. 10. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kania, T

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/EP 97/02793

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 521 621 A (CAMBRIDGE ADVANCED TECH) 7 January 1993 see the whole document ---	1-29
A	DENYER K. ET AL.: "Identification of multiple isoforms of soluble and granule-bound starch synthase in developing wheat endosperm" PLANTA, vol. 196, 1995, pages 256-265, XP002043126 cited in the application see the whole document ---	1-29
A	RAHMAN S. ET AL.: "The major proteins of wheat endosperm starch granules" AUSTRALIAN JOURNAL OF PLANT PHYSIOLOGY, vol. 22, 1995, pages 793-803, XP002043127 cited in the application see the whole document -----	1-29

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/02793

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 4441408 A	15-05-96	AU 3927995 A WO 9615248 A EP 0791066 A	06-06-96 23-05-96 27-08-97
EP 0521621 A	07-01-93	CA 2071010 A US 5365016 A	13-12-92 15-11-94

THIS PAGE BLANK (USPTO)